

AN INITIATIVE BY IITian

MODULUS
STUDY MATERIAL

TARGET YOUR JEE WITH IITians

MATHEMATICS

IIT-JEE FOUNDATION (IX and X)

www.modulushyd.com

Ph. 7 99 33 74 333 , 7 99 33 75 333 , 7 99 33 76 333

**Do you know how much percentage of marks
 a student has to score in IITJEE advanced exam to get
 a rank in top 100 ?
 a seat getting rank in IITs ?**

Unbelievably they are just 75% and 35% (avg of last 10 years data)
 Many students who score 10/10 in boards are not able to score 35% in
 JEE advanced but an average student can make it to IIT if the coaching institute has

- ✓ **Expert faculty - Excellent teaching**
- ✓ **Meticulous planning - Long-term strategy**
- ✓ **Constant motivation - Continuous support**
- ✓ **Rigorous practice - Accuracy and Speed**

Previous Years JEE ADVANCED Rank VS % of Marks

RANK	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	Avg %	2021		2021	
													(M. Marks 360)	Rank	(M. Marks 360)	Rank
1	85.48	91.67	96.01	92.22	92.78	93.06	86.02	92.62	93.61	93.01	88.89	91.40	348	1	196	1544
10	81.39	85.42	88.48	85.83	86.94	83.92	78.49	86.88	86.11	82.80	81.06	84.99	324	8	185	2043
100	70.76	79.79	78.68	76.39	77.50	69.84	68.61	83.33	74.72	72.85	71.72	74.93	323	13	178	2395
500	62.17	72.08	69.36	66.67	68.61	60.32	54.57	77.32	64.44	61.02	59.34	65.08	313	25	160	3542
1000	57.87	68.13	64.46	61.67	63.33	56.75	49.46	72.95	58.59	55.65	53.28	60.22	310	27	154	4137
2000	52.97	62.50	58.33	55.56	57.22	49.21	43.55	67.48	53.06	50.00	46.46	54.21	302	46	145	5005
3000	49.69	59.17	54.41	51.94	53.61	45.44	42.74	63.38	49.17	46.77	42.42	50.79	297	66	135	6180
4000	47.03	56.67	51.72	49.17	51.11	42.66	37.90	60.38	46.39	44.09	39.39	47.86	296	70	131	6750
5000	44.99	54.38	49.51	47.22	48.89	40.48	36.02	57.65	43.89	41.94	37.12	45.64	286	94	126	7516
6000	43.35	52.17	47.55	45.56	47.22	38.69	34.41	55.45	41.94	40.32	35.10	43.85	266	201	121	8494
7000	41.92	51.04	45.83	43.89	45.56	37.10	33.33	53.55	40.28	38.71	33.59	42.25	254	299	109	10848
8000	40.49	49.79	43.38	42.50	44.44	35.91	31.99	51.63	38.61	37.63	32.32	40.79	245	395	102	12874
9000	39.47	48.54	43.14	41.11	43.06	34.52	30.91	50.27	37.22	36.29	30.81	39.58	239	471	100	13215
10000	38.85	47.71	41.91	40.00	41.94	33.33	29.84	48.90	36.11	35.22	29.80	38.51	231	600	85	18547
QUAL%	38.85	47.71	34.55	33.88	35.00	23.81	20.16	22.50	25.00	25.00	17.42	29.44	212	1006	79	21157

**For query Call : 7993376333, 7993375333, 7993374333
 Visit : www.modulushyd.com**

1

Sequence and Series

ARITHMETIC PROGRESSION (A.P.)

KEY FACTS

1. A **sequence** is a set of numbers specified in a definite order by some assigned rule or law.

Ex. 2, 7, 12, 17.... (Each succeeding term is obtained by adding 5 to the preceding term)

1, 2, 4, 8, 16.... (Each succeeding term is obtained by multiplying the preceding term by 2)

A **finite sequence** is that which ends or has a last term.

Ex. 5, 9, 13, 17, 21.

An **infinite sequence** is one which has no last term.

Ex. 3, 6, 12, 24, 48....

In general, a_n or T_n denotes the **n th term** of a sequence.

2. An expression consisting of the term of a sequence, alternating with the symbol '+' is called a **series**.

Ex. The sequence $\frac{2}{3}, \frac{4}{5}, \frac{6}{7}, \dots$ expressed as a series is $\frac{2}{3} + \frac{4}{5} + \frac{6}{7} + \dots$

3. **Arithmetic Progression (A.P.):** A sequence is called an arithmetic progression if its terms continually increase or decrease by the same number. The fixed number by which they increase or decrease is called the **common difference**. Three quantities a, b, c will be in A.P. if $b - a = c - b$, i.e., $2b = a + c$.

(a) **n th term of an A.P.:** The n th term of an A.P. $a, a + d, a + 2d, a + 3d, \dots$ is

$$T_n = a + (n - 1)d$$

where T_n denotes n th term, a the first term, n the number of terms and d the common difference.

Also, common difference $d = T_n - T_{n-1}$.

Ex. The 9th term of the A.P.: 2, 5, 8.... is

$$T_9 = 2 + (9 - 1) \times 3 = 2 + 24 = 26.$$

Note: Here $a = 2, d = 3$.

(b) **Sum of n terms of an A.P.**

Let the A.P. be $a, a + d, a + 2d, \dots$. Let l be the last term and S the required sum. Then,

$$\begin{aligned} S &= \frac{n}{2} (a + l) = \frac{\text{Number of terms}}{2} (\text{First term} + \text{Last term}) \\ &= \frac{n}{2} (a + a + (n - 1)d) = \frac{n}{2} (2a + (n - 1)d), \end{aligned}$$

where n is the number of terms, a is first term and d is common difference.

Also, n th term = Sum of n terms - Sum of $(n - 1)$ terms

i.e., $T_n = S_n - S_{n-1}$.

Ex. The sum of 20 terms of the the A.P. 1, 3, 5, 7, 9.... is

$$S_{20} = \frac{20}{2} (2 \times 1 + (20-1) \times 2) \quad (\because a = 1, d = 2, n = 20)$$

$$= 10 \times 40 = \mathbf{400}.$$

(c) **Arithmetic Mean:** The Arithmetic Mean between two numbers is the number which when placed between them forms an arithmetic progression with them. Thus if x is the arithmetic mean of two given numbers a and b , then a, x, b form an A.P.

$$\therefore x - a = b - x \Rightarrow x = \frac{a + b}{2}$$

Ex. (a) The arithmetic mean between -4 and 6 is $\frac{6 + (-4)}{2} = \mathbf{1}$.

(b) Find 4 arithmetic means between 3 and 23.

Let A_1, A_2, A_3, A_4 , be the four arithmetic means between 3 and 23.

Then, $3, A_1, A_2, A_3, A_4, 23$ form an A.P.

Here, First term $= a = 3$

Sixth term $= T_6 = 23$

Number of terms $= n = 6$

Common difference $= d = ?$

$$T_6 = a + (n - 1) \times d$$

$$\Rightarrow 23 = 3 + (6 - 1) \times d$$

$$\Rightarrow 23 = 3 + 5d \Rightarrow 5d = 20 \Rightarrow d = \mathbf{4}.$$

$$\therefore A_1 = 3 + 4 = 7, A_2 = 7 + 4 = 11, A_3 = 11 + 4 = 15, A_4 = 15 + 4 = 19.$$

(d) Some useful facts about an A.P.

I. If each term of a given A.P. is increased or decreased or multiplied or divided by the same number, the resulting progression is also an A.P.

II. If a, b, c are in A.P., then $\frac{a - b}{b - c} = 1$, i.e., $\frac{\text{First term} - \text{Second term}}{\text{Second term} - \text{Third term}} = 1$.

(e) If we have to find an odd number of terms in A.P. whose sum is given, it is convenient to take a as the middle term and d as the common difference. Thus, three terms may be taken as $a - d, a, a + d$ and five terms as $a - 2d, a - d, a, a + d, a + 2d$. [Solved Ex. 13, 14]. If we have to find even number of terms, we take $a - d, a + d$ as the middle terms and $2d$ as the common difference. Then, four terms are taken as $a - 3d, a - d, a + d, a + 3d$.

SOLVED EXAMPLES

Ex. 1. The 8th term of a series in A.P. is 23 and the 102th term is 305. Find the series.

Sol. Let a be the first term and d be the common difference.

$$\text{Then, } T_8 = a + (8 - 1)d \Rightarrow 23 = a + 7d \quad \dots(i)$$

$$T_{102} = a + (102 - 1)d \Rightarrow 305 = a + 101d \quad \dots(ii)$$

$$\therefore \text{Eqn (ii)} - \text{Eqn (i)}$$

$$\Rightarrow 94d = 282 \quad \Rightarrow d = 3$$

Now substituting $d = 3$ in (i), we get $23 = a + 21 \Rightarrow a = 2$.

$$\therefore a = 2, d = 3 \Rightarrow \text{Series is } 2 + 5 + 8 + 11 + \dots$$

Ex. 2. If a, b and c be respectively the p th, q th and r th terms of an A.P., prove that $a(q - r) + b(r - p) + c(p - q) = 0$.

Sol. Let A be the first term and D the common difference of the given A.P.

$$\text{Then, } T_p = A + (p - 1)D = a \quad \dots(i)$$

$$T_q = A + (q - 1)D = b \quad \dots(ii)$$

$$T_r = A + (r - 1)D = c \quad \dots(iii)$$

$$\begin{aligned}
&\therefore (i) \times (q-r) + (ii) \times (r-p) + (iii) \times (p-q) \\
&\Rightarrow A [(q-r) + (r-p) + (p-q)] + D [(p-1)(q-r) + (q-1)(r-p) + (r-1)(p-q)] \\
&\hspace{20em} = a(q-r) + b(r-p) + c(p-q) \\
&\Rightarrow A [0] + D [pq - q - pr + r + qr - r - pq + p + pr - p - qr + q] = a(q-r) + b(r-p) + c(p-q) \\
&\Rightarrow a(q-r) + b(r-p) + c(p-q) = A \times 0 + D \times 0 = 0.
\end{aligned}$$

Ex. 3. Which term of the progression $19, 18\frac{1}{5}, 17\frac{2}{5}, \dots$ is the first negative term?

Sol. Here $a = 19, d = 18\frac{1}{5} - 19 = -\frac{4}{5}$

Let the n th term be the first negative term. Then,

$$\begin{aligned}
T_n < 0 &\Rightarrow a + (n-1)d < 0 \\
&\Rightarrow 19 + (n-1)\left(-\frac{4}{5}\right) < 0 \\
&\Rightarrow 19 - \frac{4}{5}n + \frac{4}{5} < 0 \\
&\Rightarrow \frac{99}{5} - \frac{4}{5}n < 0 \Rightarrow n > \frac{99}{5} \times \frac{5}{4} \Rightarrow n > \frac{99}{4} = 24\frac{3}{4} \\
&\Rightarrow n = 25.
\end{aligned}$$

Ex. 4. Find the sum of the series $101 + 99 + 97 + \dots + 47$.

Sol. In this case, we have to first find the number of terms.

Here $a = 101, l = T_n = 47, d = 99 - 101 = -2$

$\therefore 47 = 101 + (n-1) \times (-2)$, where $n =$ number of terms

$\Rightarrow 47 = 101 - 2n + 2$

$\Rightarrow 2n = 103 - 47 = 56 \Rightarrow n = 28$

$\therefore S_n = \frac{n}{2}(a+l)$

$\Rightarrow S_{28} = \frac{28}{2}(101+47) = 14 \times 148 = 2072.$

Ex. 5. The sums of n terms of two arithmetic series are in the ratio $2n+1 : 2n-1$. Find the ratio of their 10th terms.

Sol. Let the two arithmetic series be $a, a+d, a+2d, \dots$ and $A, A+D, A+2D, \dots$

Given that, $\frac{n/2[2a+(n-1)d]}{n/2[2A+(n-1)D]} = \frac{2n+1}{2n-1}$

$\Rightarrow \frac{2a+(n-1)d}{2A+(n-1)D} = \frac{2n+1}{2n-1}$...(i)

Ratio of the 10th terms of these series $= \frac{t_{10}}{T_{10}} = \frac{a+9d}{A+9D} = \frac{2a+18d}{2A+18D}$

\therefore Putting $n = 19$ in (i), we have $\frac{t_{10}}{T_{10}} = \frac{2a+18d}{2A+18D} = \frac{2 \times 19 + 1}{2 \times 19 - 1} = \frac{39}{37}.$

Ex. 6. The sum of the first n terms of the arithmetical progression $3, 5\frac{1}{2}, 8, \dots$ is equal to the $2n$ th term of the A.P. $16\frac{1}{2}, 28\frac{1}{2}, 40\frac{1}{2}, \dots$. Calculate the value of n .

Sol. For the A.P. : $3, 5\frac{1}{2}, 8, \dots$ $a = 3, d = 2\frac{1}{2}$, number of terms = n

$$\Rightarrow S_n = \frac{n}{2} [2a + (n-1)d] = \frac{n}{2} \left(6 + (n-1) \times \frac{5}{2} \right) \quad \dots(i)$$

For the A.P.: $16\frac{1}{2}, 28\frac{1}{2}, 40\frac{1}{2} \dots$ $a = 16\frac{1}{2}, d = 12$.

$$\therefore T_{2n} = a + (2n-1)d = 16\frac{1}{2} + (2n-1) \times 12 \quad \dots(ii)$$

$$\text{Given, } S_n = T_{2n} \Rightarrow \frac{n}{2} \left(6 + (n-1) \frac{5}{2} \right) = \frac{33}{2} + (2n-1) \times 12 \quad (\text{From (i) and (ii)})$$

$$\Rightarrow \frac{6n}{2} + \frac{5n^2}{4} - \frac{5n}{4} = \frac{33}{2} + 24n - 12 \quad \Rightarrow \frac{7n}{4} + \frac{5n^2}{4} = \frac{9}{2} + 24n$$

$$\Rightarrow \frac{5n^2}{4} - \frac{89n}{4} - \frac{9}{2} = 0 \quad \Rightarrow 5n^2 - 89n - 18 = 0$$

$$\Rightarrow 5n^2 - 90n + n - 18 = 0 \quad \Rightarrow 5n(n-18) + 1(n-18) = 0$$

$$\Rightarrow (n-18)(5n+1) = 0 \quad \Rightarrow n = 18 \text{ or } -\frac{1}{5}$$

\Rightarrow Neglecting $-ve$ value, we have $n = 18$.

Ex. 7. Let a_1, a_2, a_3, \dots be the terms of an A.P. If $\frac{a_1 + a_2 + a_3 + \dots + a_p}{a_1 + a_2 + a_3 + \dots + a_q} = \frac{p^2}{q^2}$ ($p \neq q$), then find $\frac{a_6}{a_{21}}$.

(AIEEE 2006)

Sol. Let d be the common difference for the A.P.; a_1, a_2, a_3, \dots

$$\text{Then, } \frac{S_p}{S_q} = \frac{p/2 [2a_1 + (p-1)d]}{q/2 [2a_1 + (q-1)d]} = \frac{p^2}{q^2}$$

$$\Rightarrow \frac{2a_1 + (p-1)d}{2a_1 + (q-1)d} = \frac{p}{q} \quad \Rightarrow q [2a_1 + (p-1)d] = p [2a_1 + (q-1)d]$$

$$\Rightarrow 2a_1q + pqd - qd = 2a_1p + pqd - pd \quad \Rightarrow pd - qd = 2a_1p - 2a_1q$$

$$\Rightarrow d(p-q) = 2a_1(p-q) \quad \Rightarrow d = 2a_1$$

$$\text{Now } \frac{\text{Term 6}}{\text{Term 21}} = \frac{a_6}{a_{21}} = \frac{a_1 + (6-1)d}{a_1 + (21-1)d} = \frac{a_1 + 5 \times 2a_1}{a_1 + 20 \times 2a_1} = \frac{11a_1}{41a_1} = \frac{11}{41}$$

Ex. 8. If the number of terms of an A.P. is $(2n+1)$, then what is the ratio of the sum of the odd terms to the sum of even terms? (NDA/NA 2008)

Sol. Let the A.P. be $a, a+d, a+2d, \dots, a+(2n-1)d, a+2nd$

Then, the progression of odd terms is $a, a+2d, a+4d, \dots, a+2nd$.

This progression has $(n+1)$ terms.

$$\text{Its sum} = \frac{n+1}{2} [a + a + 2nd] = \frac{n+1}{2} [2a + 2nd] = (n+1)(a + nd) \quad \dots(i)$$

The progression of even terms is $a+d, a+3d, \dots, a+(2n-1)d$.

This progression has n terms.

$$\text{Its sum} = \frac{n}{2} [(a + d) + (a + (2n - 1)d)] = \frac{n}{2} [2a + 2nd] = n(a + nd) \quad \dots(ii)$$

$$\therefore \frac{\text{Sum of odd terms}}{\text{Sum of even terms}} = \frac{(n + 1)(a + nd)}{n(a + nd)} = \frac{n + 1}{n}$$

Ex. 9. If the sum of the roots of the quadratic equation $ax^2 + bx + c = 0$ is equal to the sum of the squares of their reciprocals, then show that ab^2, ca^2, bc^2 are in A.P. (DCE)

Sol. Let α, β be the roots of the equation $ax^2 + bx + c = 0$.

Then $\alpha + \beta = -b/a, \alpha\beta = c/a$

Given, Sum of roots = Sum of squares of reciprocals of roots

$$\Rightarrow \alpha + \beta = \frac{1}{\alpha^2} + \frac{1}{\beta^2} \quad \Rightarrow \alpha + \beta = \frac{\alpha^2 + \beta^2}{\alpha^2\beta^2} \quad \Rightarrow \alpha + \beta = \frac{(\alpha + \beta)^2 - 2\alpha\beta}{(\alpha\beta)^2}$$

$$\Rightarrow -b/a = \frac{(-b/a)^2 - \frac{2c}{a}}{(c/a)^2} \quad \Rightarrow \frac{-b}{a} = \frac{b^2 - 2ca}{c^2} \quad \Rightarrow -bc^2 = ab^2 - 2ca^2$$

$$\Rightarrow 2ca^2 = ab^2 + bc^2 \quad \Rightarrow ab^2, ca^2, bc^2 \text{ are in A.P.} \quad (\because a, b, c \text{ in A.P.} \Rightarrow 2b = a + c)$$

Ex. 10. Find the value of n , if $\frac{a^{n+1} + b^{n+1}}{a^n + b^n}$ is the arithmetic mean between a and b . (WBJEE 2009)

Sol. The arithmetic mean between a and b is $\frac{a + b}{2}$.

Given, $\frac{a + b}{2} = \frac{a^{n+1} + b^{n+1}}{a^n + b^n}$

$$\Rightarrow (a^n + b^n)(a + b) = 2(a^{n+1} + b^{n+1}) \quad \Rightarrow a^{n+1} + a^n b + b^n a + b^{n+1} = 2a^{n+1} + 2b^{n+1}$$

$$\Rightarrow a^{n+1} + b^{n+1} = a^n b + b^n a \quad \Rightarrow (a^{n+1} - b a^n) - (b^n a - b^{n+1}) = 0$$

$$\Rightarrow a^n(a - b) - b^n(a - b) = 0 \quad \Rightarrow (a^n - b^n)(a - b) = 0$$

$$\Rightarrow (a^n - b^n) = 0 \text{ or } (a - b) = 0$$

But $a \neq b \Rightarrow a - b \neq 0$

$$\therefore a^n - b^n = 0 \Rightarrow a^n = b^n \Rightarrow \left(\frac{a}{b}\right)^n = 1 \Rightarrow \left(\frac{a}{b}\right)^n = \left(\frac{a}{b}\right)^0 \Rightarrow n = 0.$$

Ex. 11. If $\log_{10} 2, \log_{10} (2^x - 1)$ and $\log_{10} (2^x + 3)$ be three consecutive terms of an A.P., then find the value of x . (AMU 2012)

Sol. Given, $\log_{10} 2, \log_{10} (2^x - 1), \log_{10} (2^x + 3)$ are in A.P. Then,

$$\log_{10} (2^x - 1) - \log_{10} 2 = \log_{10} (2^x + 3) - \log_{10} (2^x - 1)$$

$$\Rightarrow \log_{10} \left(\frac{2^x - 1}{2}\right) = \log_{10} \left(\frac{2^x + 3}{2^x - 1}\right)$$

$$\Rightarrow \frac{2^x - 1}{2} = \frac{2^x + 3}{2^x - 1} \quad \Rightarrow (2^x - 1)^2 = 2(2^x + 3)$$

$$\Rightarrow 2^{2x} - 2 \cdot 2^x + 1 = 2 \cdot 2^x + 6 \quad \Rightarrow 2^{2x} - 4 \cdot 2^x - 5 = 0$$

$$\Rightarrow 2^{2x} - 5 \cdot 2^x + 2^x - 5 = 0 \quad \Rightarrow 2^x(2^x - 5) + 1(2^x - 5) = 0$$

$$\Rightarrow (2^x - 5)(2^x + 1) = 0 \quad \Rightarrow 2^x = 5 \quad (\because 2^x \neq -1)$$

$$\Rightarrow x = \log_2 5.$$

Ex. 12. If $a_1, a_2, a_3, \dots, a_n$ be an A.P. of non-zero terms, then find the sum: $\frac{1}{a_1 a_2} + \frac{1}{a_2 a_3} + \dots + \frac{1}{a_{n-1} a_n}$. (AMU 2009)

Sol. Let $a_2 - a_1 = a_3 - a_2 = \dots = a_n - a_{n-1} = d$ (common difference)

$$\begin{aligned} \text{Then, } \frac{1}{a_1 a_2} + \frac{1}{a_2 a_3} + \dots + \frac{1}{a_{n-1} a_n} &= \frac{1}{d} \left[\frac{d}{a_1 a_2} + \frac{d}{a_2 a_3} + \dots + \frac{d}{a_{n-1} a_n} \right] \\ &= \frac{1}{d} \left[\frac{a_2 - a_1}{a_1 a_2} + \frac{a_3 - a_2}{a_2 a_3} + \dots + \frac{a_n - a_{n-1}}{a_{n-1} a_n} \right] = \frac{1}{d} \left[\frac{1}{a_1} - \frac{1}{a_2} + \frac{1}{a_2} - \frac{1}{a_3} + \dots + \frac{1}{a_{n-1}} - \frac{1}{a_n} \right] \\ &= \frac{1}{d} \left[\frac{1}{a_1} - \frac{1}{a_n} \right] = \frac{1}{d} \left[\frac{a_n - a_1}{a_1 a_n} \right] = \frac{1}{d} \left[\frac{(a_1 + (n-1)d) - a_1}{a_1 a_n} \right] = \frac{n-1}{a_1 a_n}. \end{aligned}$$

Ex. 13. If the sides of a right angled triangle form an A.P., then find the sines of the acute angles. (VITEE 2008)

Sol. Let the $\triangle ABC$ be right angled at C.

Then, $AB = c, BC = a, AC = b$

Given, the sides of the right angled \triangle are in A.P. $\Rightarrow a, b, c$ are in A.P.

Now, let $a = x - d, b = x, c = x + d$

(d being a +ve quantity as c being the hypotenuse is the greatest side)

$$\therefore c^2 = a^2 + b^2 \quad (\text{Pythagoras' Theorem})$$

$$\Rightarrow (x + d)^2 = (x - d)^2 + x^2$$

$$\Rightarrow x^2 + 2xd + d^2 = x^2 - 2xd + d^2 + x^2$$


$$\Rightarrow 4xd = x^2 \quad \Rightarrow \quad d = \frac{x}{4}.$$

$$\therefore a = x - d = x - \frac{x}{4} = \frac{3x}{4}, \quad b = x, \quad c = x + d = x + \frac{x}{4} = \frac{5x}{4}$$

C being the right angle, A and B are the acute angles.

$$\therefore \sin A = \frac{a}{c} = \frac{3x/4}{5x/4} = \frac{3}{5} \quad \text{and} \quad \sin B = \frac{b}{c} = \frac{x/5}{x/4} = \frac{x}{5x} = \frac{4}{5}$$

\therefore The sines of the acute angles are $\frac{3}{5}, \frac{4}{5}$.



Ex. 14. a_1, a_2, a_3, a_4, a_5 are the first five terms of an A.P. such that $a_1 + a_3 + a_5 = -12$ and $a_1 \cdot a_2 \cdot a_3 = 8$. Find the first term and common difference.

Sol. Let $a_1 = a_3 - 2d, a_2 = a_3 - d, a_3 = a_3, a_4 = a_3 + d, a_5 = a_3 + 2d$

$$\text{Then } a_1 + a_3 + a_5 = -12, \text{ (given)} \Rightarrow a_3 - 2d + a_3 + a_3 + 2d = -12 \Rightarrow 3a_3 = -12 \Rightarrow a_3 = -4$$

Also, $a_1 \cdot a_2 \cdot a_3 = 8$ (given)

$$\Rightarrow a_1 \cdot a_2 = -2 \quad (\because a_3 = -4) \Rightarrow (a_3 - 2d)(a_3 - d) = -2$$

$$\Rightarrow (-4 - 2d)(-4 - d) = -2 \Rightarrow (2 + d)(4 + d) = 1$$

$$\Rightarrow d^2 + 6d + 9 = 0 \quad \Rightarrow \quad (d + 3)^2 = 0 \Rightarrow d = -3$$

$$\therefore a_1 = a_3 - 2d = -4 - 2(-3) = 2.$$

Ex. 15. If a, b, c are in A.P. show that $\frac{a(b+c)}{bc}, \frac{b(c+a)}{ca}, \frac{c(a+b)}{ab}$ are in A.P.

Sol. a, b, c are in A.P.

$$\Rightarrow \frac{a}{abc}, \frac{b}{abc}, \frac{c}{abc} \text{ are in A.P.}$$

(Dividing each term by abc)

$$\Rightarrow \frac{1}{bc}, \frac{1}{ac}, \frac{1}{ab} \text{ are in A.P.}$$

$$\Rightarrow \frac{ab+bc+ca}{bc}, \frac{ab+bc+ca}{ac}, \frac{ab+bc+ca}{ab} \text{ are in A.P.} \quad (\text{Multiplying each term by } ab+bc+ca)$$

$$\Rightarrow \frac{ab+ca}{bc} + 1, \frac{ab+bc}{ac} + 1, \frac{bc+ca}{ab} + 1 \text{ are in A.P.}$$

$$\Rightarrow \frac{ab+ca}{bc}, \frac{ab+bc}{ac}, \frac{bc+ca}{ab} \text{ are in A.P.} \quad (\text{Subtracting 1 from each term})$$

$$\Rightarrow \frac{a(b+c)}{bc}, \frac{b(a+c)}{ac}, \frac{c(b+a)}{ab} \text{ are in A.P.}$$

Ex. 16. If $\frac{b+c-a}{a}, \frac{c+a-b}{b}, \frac{a+b-c}{c}$ are in A.P., prove that $\frac{1}{a}, \frac{1}{b}, \frac{1}{c}$ are also in A.P.

Sol. $\frac{b+c-a}{a}, \frac{c+a-b}{b}, \frac{a+b-c}{c}$ are in A.P.

$$\Rightarrow \frac{b+c-a}{a} + 2, \frac{c+a-b}{b} + 2, \frac{a+b-c}{c} + 2 \text{ are in A.P.} \quad (\text{Adding 2 to each term of A.P.})$$

$$\Rightarrow \frac{b+c+a}{a}, \frac{c+a+b}{b}, \frac{a+b+c}{c} \text{ are in A.P.}$$

$$\Rightarrow \frac{1}{a}, \frac{1}{b}, \frac{1}{c} \text{ are in A.P.} \quad (\text{Dividing each term by } a+b+c)$$

Ex. 17. If a, b, c are in A.P., show that $\frac{1}{\sqrt{b}+\sqrt{c}}, \frac{1}{\sqrt{c}+\sqrt{a}}, \frac{1}{\sqrt{a}+\sqrt{b}}$ are in A.P. (NDA/NA 2010)

Sol. Given a, b, c are in A.P. $\Rightarrow 2b = a + c$

Now $\frac{1}{\sqrt{b}+\sqrt{c}}, \frac{1}{\sqrt{c}+\sqrt{a}}, \frac{1}{\sqrt{a}+\sqrt{b}}$ are in A.P., if

$$\frac{1}{\sqrt{c}+\sqrt{a}} - \frac{1}{\sqrt{b}+\sqrt{c}} = \frac{1}{\sqrt{a}+\sqrt{b}} - \frac{1}{\sqrt{c}+\sqrt{a}}$$

$$\Rightarrow \frac{2}{\sqrt{c}+\sqrt{a}} = \frac{1}{\sqrt{b}+\sqrt{c}} + \frac{1}{\sqrt{a}+\sqrt{b}} \Rightarrow \frac{2}{\sqrt{c}+\sqrt{a}} = \frac{\sqrt{a}+\sqrt{b}+\sqrt{b}+\sqrt{c}}{(\sqrt{b}+\sqrt{c})(\sqrt{a}+\sqrt{b})}$$

$$\Rightarrow 2(\sqrt{b}+\sqrt{c})(\sqrt{a}+\sqrt{b}) = (\sqrt{c}+\sqrt{a})(\sqrt{a}+2\sqrt{b}+\sqrt{c})$$

$$\Rightarrow 2(\sqrt{ab}+b+\sqrt{ac}+\sqrt{bc}) = (\sqrt{ac}+2\sqrt{bc}+c+a+2\sqrt{ba}+\sqrt{ac})$$

$$\Rightarrow 2\sqrt{ab}+2b+2\sqrt{ac}+2\sqrt{bc} = 2\sqrt{ac}+2\sqrt{bc}+c+a+2\sqrt{ba}$$

$$\Rightarrow 2b = a + c, \text{ which is true as } a, b, c \text{ are in A.P.}$$

$$\Rightarrow \frac{1}{\sqrt{b}+\sqrt{c}}, \frac{1}{\sqrt{c}+\sqrt{a}}, \frac{1}{\sqrt{a}+\sqrt{b}} \text{ are in A.P.}$$

Ex. 18. What is the sum of all two-digit numbers which leave remainder 5 when they are divided by 7?

Sol. The two digit natural numbers which leave a remainder 5, when divided by 7 are 12, 19, 26, ..., 89, 96.

\therefore 12, 19, 26, ..., 89, 96 is an A.P. whose first term $a = 12$ and common difference $d = 7$.

Let the last or n th term be T_n .

Then, $T_n = a + (n-1)d$, where n is the number of terms in A.P.

$$\Rightarrow 96 = 12 + (n-1)7$$

$$\Rightarrow 84 = (n-1)7 \Rightarrow n-1 = 12 \Rightarrow n = 13$$

$$\therefore \text{Required Sum} = \frac{n}{2}(a+l) = \frac{13}{2}(12+96) = \frac{13}{2} \times 108 = 13 \times 54 = 702.$$

Ex. 19. What is the sum of the series $1^2 - 2^2 + 3^2 - 4^2 + \dots + 99^2 - 100^2$.

(Orissa JEE 2006)

Sol. $1^2 - 2^2 + 3^2 - 4^2 + 5^2 - 6^2 + \dots + 99^2 - 100^2$

$$= (1-2)(1+2) + (3-4)(3+4) + (5-6)(5+6) + \dots + (99-100)(99+100)$$

$$= -(1+2) - (3+4) - (5+6) \dots - (99+100)$$

$$= -(1+2+3+4+5+6+\dots+99+100) = -\left(\frac{100}{2}(1+100)\right) \quad (\because S_n = \frac{n}{2}(\text{first term} + \text{last term}))$$

$$= -50 \times 101 = -5050.$$

Ex. 20. If the first, second and last terms of an arithmetic series are a , b and c respectively, then what is the number of terms?

(MPPET 2009)

Sol. $\left. \begin{array}{l} \text{First term} = a \\ \text{Second term} = b \end{array} \right\} \Rightarrow \text{common difference } (d) = b - a$

Last term = c

Let the number of terms be n . Then,

$$c = a + (n-1)d \Rightarrow c = a + (n-1)(b-a) \Rightarrow c - a = (n-1)(b-a)$$

$$\Rightarrow (n-1) = \frac{c-a}{b-a} \Rightarrow n = \frac{c-a}{b-a} + 1 = \frac{c-a+b-a}{b-a} = \frac{c+b-2a}{b-a}.$$

Ex. 21. After inserting x A.M's. between 2 and 38, the sum of the resulting progression is 200. What is the value of x ?

(AMU 2001)

Sol. After inserting x A.M's between 2 and 38, we get an A.P. of $(x+2)$ terms with first term as 2 and last term as 38.

Now, sum of n terms of an A.P. = $\frac{n}{2}(a+l)$, where a = first term, l = last term

$$\therefore \text{Here, } 200 = \frac{(x+2)}{2}(2+38) \Rightarrow 200 = 20(x+2)$$

$$\Rightarrow 20x + 40 = 200 \Rightarrow 20x = 160 \Rightarrow x = 8.$$

PRACTICE SHEET

1. Find the 26th term of the A.P: 10, 6, 2, -2, -6, -10, ?

(a) -86 (b) 96 (c) -90 (d) -106

2. The 2nd, 31st and last term of an A.P. are $7\frac{3}{4}$, $\frac{1}{2}$ and $-6\frac{1}{2}$

respectively. The number of terms of the A.P. is

(a) 48 (b) 60 (c) 52 (d) 59

3. If p times the p th term of an A.P. is q times the q th term, then what is $(p+q)$ th term equal to?

(a) $p+q$ (b) pq (c) 1 (d) 0

(NDA/NA 2010)

4. The 59th term of an A.P. is 449 and the 449th term is 59. Which term is equal to 0?

(a) 501st term (b) 502nd term
(c) 508th term (d) 509th term

(NDA/NA 2010)

5. Let T_r be the r th term of an A.P. whose first term is a and common difference is d . If for some positive integers m and

n , $T_m = \frac{1}{n}$ and $T_n = \frac{1}{m}$, then $(a-d)$ equals

(a) $1/mn$ (b) 1 (c) 0 (d) $\frac{1}{m} + \frac{1}{n}$

(AIEEE 2004, UPSEE 2007)

6. If a, b, c be in Arithmetic Progression, then the value of $(a+2b-c)(2b+c-a)(a+2b+c)$ is

(a) $3abc$ (b) $4abc$ (c) $8abc$ (d) $16abc$

(WBEE 2008)

7. Find the sum of 24 terms of the series $2\frac{1}{2}, 3\frac{1}{3}, 4\frac{1}{6}, 5, \dots$?

(a) 200 (b) 185 (c) 290 (d) 250

8. If the sum of the 12th and 22nd terms of an A.P. is 100, then the sum of the first 33 terms of the A.P. is

(a) 1650 (b) 2340 (c) 3300 (d) 3400

(Kerala PET 2008)

9. If $S_1 = a_2 + a_4 + a_6 + \dots$ upto 100 terms and $S_2 = a_1 + a_3 + a_5 + \dots$ upto 100 terms of a certain A.P., then its common difference is
 (a) $S_1 - S_2$ (b) $S_2 - S_1$
 (c) $\frac{S_1 - S_2}{2}$ (d) None of these
 (AMU 2010)
10. The sum of n terms of an A.P. is $2n + 3n^2$. Which term of this A.P. is equal to 299?
 (a) 11th (b) 50th (c) 35th (d) 29th
11. If S_n denotes the sum of first n terms of an A.P. $a_1 + a_2 + a_3 + \dots$, such that $\frac{S_m}{S_n} = \frac{m^2}{n^2}$, then $\frac{a_m}{a_n} = ?$
 (a) $m - 1 : n - 1$ (b) $m - n : m + n$
 (c) $2m - 1 : 2n - 1$ (d) $m + 1 : n + 1$
 (J&K CET 2013)
12. If the sum of the first ten terms of an A.P. is 4 times the sum of the first five terms, then the ratio of the first term to the common difference is:
 (a) 1 : 2 (b) 2 : 1 (c) 1 : 4 (d) 4 : 1
 (NDA/NA 2003)
13. What is the sum of numbers lying between 107 and 253, which are divisible by 5?
 (a) 5250 (b) 5210 (c) 5220 (d) 5000
14. The ratio between the sum of n terms of two A.P.'s is $(7n + 1) : (4n + 27)$. The ratio of their 11th terms is
 (a) 125 : 106 (b) 148 : 111 (c) 131 : 89 (d) 127 : 108
15. Let S_n denote the sum of first n terms of an A.P. If $S_{2n} = 3S_n$, then the ratio $\frac{S_{3n}}{S_n}$ is equal to
 (a) 4 (b) 6 (c) 8 (d) 10
 (MAT 2002, Rajasthan PET 2006)
16. If the sum of $2n$ terms of the A.P. 2, 5, 8, 11, ... is equal to the sum of n terms of A.P. 57, 59, 61, 63, ..., then n is equal to
 (a) 10 (b) 11 (c) 12 (d) 13
 (IIT 2001)
17. If S_1, S_2, S_3 denote respectively the sum of first n_1, n_2 and n_3 terms of an A.P., then
 $\frac{S_1}{n_1}(n_2 - n_3) + \frac{S_2}{n_2}(n_3 - n_1) + \frac{S_3}{n_3}(n_1 - n_2)$ is equal to
 (a) 0 (b) $n_1 + n_2 + n_3$
 (c) $n_1 n_2 n_3$ (d) $S_1 S_2 S_3$ (DCE 2007)
18. If $S_n = nP + \frac{n}{2}(n-1)Q$, where S_n denotes the sum of first n terms of an A.P., then the common difference of the A.P. is
 (a) Q (b) $P + Q$ (c) $P + 2Q$ (d) $2P + 3Q$
 (DCE 2001)
19. n arithmetic means are inserted between 3 and 17. If the ratio of the last and the first arithmetic mean is 3 : 1, then n is equal to
 (a) 5 (b) 6 (c) 7 (d) 9
 (DEC 2008)
20. If $a_1, a_2, a_3, \dots, a_n$ are in A.P. and $a_1 = 0$, then the value of $\left(\frac{a_3}{a_2} + \frac{a_4}{a_3} + \dots + \frac{a_n}{a_{n-1}}\right) - a_2 \left(\frac{1}{a_2} + \frac{1}{a_3} + \dots + \frac{1}{a_{n-2}}\right)$ is equal to
 (a) $n + \frac{1}{n}$ (b) $n + \frac{1}{n-1}$
 (c) $(n-1) + \frac{1}{(n-1)}$ (d) $(n-2) + \frac{1}{(n-2)}$
 (Kerala PET 2011)
21. Let a_1, a_2, a_3, a_4 be in A.P. If $a_1 + a_4 = 10$ and $a_2 a_3 = 24$, then the least term of them is
 (a) 1 (b) 2 (c) 3 (d) 4
 (Kerala PET 2013)
22. An A.P. has a property that the sum of first ten terms is half the sum of next ten terms. If the second term is 13, then the common difference is
 (a) 3 (b) 2 (c) 5 (d) 4
 (Kerala PET 2013)
23. The sides of a triangle are in A.P. and its area is $\frac{3}{5}$ th the area of an equilateral triangle of same perimeter. The sides of the triangle are in the ratio.
 (a) 1 : 2 : $\sqrt{7}$ (b) 2 : 3 : 5 (c) 1 : 6 : 7 (d) 3 : 5 : 7
24. If $\log_3 2, \log_3(2^x - 5)$ and $\log_3(2^x - 7/2)$ are in A.P., the value of x is
 (a) 0 (b) $\frac{1}{3}$ (c) 2 (d) 3
 (IIT 1990)
25. If $a_1, a_2, a_3, \dots, a_n$ are in A.P with common difference $d \neq 0$, then the value of $\sin d (\operatorname{cosec} a_1 \operatorname{cosec} a_2 + \operatorname{cosec} a_2 \operatorname{cosec} a_3 + \dots + \operatorname{cosec} a_{n-1} \operatorname{cosec} a_n)$ will be
 (a) $\sec a_1 - \sec a_n$ (b) $\tan a_1 - \tan a_n$
 (c) $\operatorname{cosec} a_1 - \operatorname{cosec} a_n$ (d) $\cot a_1 - \cot a_n$
 (Rajasthan PET 2000)

ANSWERS

1. (c) 2. (d) 3. (d) 4. (c) 5. (c) 6. (d) 7. (c) 8. (a) 9. (d) 10. (b)
 11. (c) 12. (a) 13. (c) 14. (b) 15. (b) 16. (b) 17. (a) 18. (a) 19. (b) 20. (d)
 21. (b) 22. (b) 23. (d) 24. (d) 25. (d)

HINTS AND SOLUTIONS

1. $T_n = a + (n-1)d \Rightarrow T_{26} = a + 25d$

Here $a = 10, d = -4$

$\therefore T_{26} = 10 + 25 \times (-4) = 10 - 100 = -90.$

2. Let the first term, common difference and number of terms of the A.P. be a, d and n respectively. Then,

$$a + d = 7\frac{3}{4} \quad \dots(i)$$

$$a + 30d = \frac{1}{2} \quad \dots(ii)$$

and $a + (n-1)d = -6\frac{1}{2} \quad \dots(iii)$

Eqn (ii) – Eqn (i)

$$\Rightarrow 29d = \frac{1}{2} - \frac{31}{4} = -\frac{29}{4}$$

$$\Rightarrow d = -\frac{1}{4}$$

Putting $d = -\frac{1}{4}$ in (i), we get

$$a - \frac{1}{4} = 7\frac{3}{4} \Rightarrow a = 7\frac{3}{4} + \frac{1}{4}$$

$$\Rightarrow a = 8$$

\therefore Putting the values of a and d in (iii), we have

$$8 + (n-1)\left(-\frac{1}{4}\right) = -\frac{13}{2}$$

$$\Rightarrow 8 + \frac{1}{4} - \frac{1}{4}n = -\frac{13}{2}$$

$$\Rightarrow -\frac{1}{4}n = -\frac{13}{2} - \frac{33}{4}$$

$$\Rightarrow -\frac{1}{4}n = -\frac{59}{4}$$

$$\Rightarrow n = 59.$$

3. Given, $p(a + (p-1)d) = q(a + (q-1)d)$, where a and d are the first term and common difference of the A.P.

$$\Rightarrow (p-q)a = (q^2 - q - p^2 + p)d$$

$$\Rightarrow -(q-p)a = (q-p)((q+p)-1)d$$

$$\Rightarrow -a = ((q+p)-1)d$$

$$\Rightarrow a + ((p+q)-1)d = 0$$

$$\Rightarrow t_{p+q} = 0.$$

4. Let a and d be the first term and common difference of the given A.P.

Then, $a + 58d = 449 \quad \dots(i)$

$a + 448d = 59 \quad \dots(ii)$

Solving eqns (i) and (ii) simultaneously, we get

$$a = 507, d = -1$$

Now assume that the n th term is zero.

$$\therefore 0 = a + (n-1)d$$

$$\Rightarrow 0 = 507 + (n-1)(-1)$$

$$\Rightarrow 507 = n-1$$

$$\Rightarrow n = 508.$$

5. Given, $T_m = \frac{1}{n} \Rightarrow a + (m-1)d = \frac{1}{n} \quad \dots(i)$

$$T_n = \frac{1}{m} \Rightarrow a + (n-1)d = \frac{1}{m} \quad \dots(ii)$$

$$\text{Eq. (ii) - Eq. (i)} \Rightarrow (n-m)d = \frac{1}{m} - \frac{1}{n}$$

$$\Rightarrow (n-m)d = \frac{n-m}{mn} \Rightarrow d = \frac{1}{mn}$$

Putting $d = \frac{1}{mn}$ in (i), we get $a = \frac{1}{mn}$

$$\therefore a - d = \frac{1}{mn} - \frac{1}{mn} = 0.$$

6. a, b, c are in A.P. $\Rightarrow 2b = a + c$

$$\therefore (a + 2b - c)(2b + c - a)(a + 2b + c)$$

$$= (a + a + c - c)(a + c + c - a)(2b + 2b)$$

$$= 2a \cdot 2c \cdot 4b = 16abc.$$

7. Here $a = \frac{5}{2}, d = \frac{10}{3} - \frac{5}{2} = \frac{20-15}{6} = \frac{5}{6}$

$$\therefore S_{24} = \frac{24}{2} \left[2 \times \frac{5}{2} + (24-1) \frac{5}{6} \right]$$

$$\left[\because S_n = \frac{n}{2} [2a + (n-1)d] \right]$$

$$= 12 \left[5 + \frac{23 \times 5}{6} \right] = 2 [30 + 115] = 2 \times 145 = 290.$$

8. Let the first term of the A.P. be a and let the common difference be d .

Then, $t_{12} + t_{22} = 100 \Rightarrow (a + 11d) + (a + 21d) = 100$
 $\Rightarrow 2a + 32d = 100 \quad \dots(i)$

Now sum of first 33 terms of the A.P

$$= \frac{33}{2} (2a + 32d) \left(\because S_n = \frac{n}{2} (2a + (n-1)d) \right)$$

$$= \frac{33}{2} \times 100 \quad \text{(From (i))}$$

$$= 1650.$$

9. Given that,

$$S_1 = a_2 + a_4 + a_6 + \dots + \text{upto 100 terms} \quad \dots(i)$$

$$S_2 = a_1 + a_3 + a_5 + \dots + \text{upto 100 terms} \quad \dots(ii)$$

Let d be the common difference of the given A.P. Then,

$$d = a_2 - a_1 = a_3 - a_2 = a_4 - a_3 = \dots \quad \dots(iii)$$

Subtracting eqn (ii) from eqn (i), we have

$$S_1 - S_2 = (a_1 - a_2) + (a_4 - a_3) + (a_6 - a_5) + \dots + \text{upto 100 terms}$$

$$= d + d + d + \dots + \text{upto 100 terms}$$

$$= 100d$$

$$\Rightarrow d = \frac{S_1 - S_2}{100}.$$

10. $S_n = 3n^2 + 2n$
 $\therefore S_{n-1} = 3(n-1)^2 + 2(n-1)$
 $= 3n^2 - 6n + 3 + 2n - 2 = 3n^2 - 4n + 1$
 $\therefore T_n = S_n - S_{n-1}$
 $= (3n^2 + 2n) - (3n^2 - 4n + 1) = 6n - 1$
 Given, $T_n = 299$
 $\Rightarrow 6n - 1 = 299$
 $\Rightarrow 6n = 300 \Rightarrow n = 50$.

11. $\frac{S_m}{S_n} = \frac{m/2 [2a_1 + (m-1)d]}{n/2 [2a_1 + (n-1)d]} = \frac{m^2}{n^2}$
 $(d \rightarrow \text{common difference of A.P.})$
 $\Rightarrow n[2a_1 + (m-1)d] = m[2a_1 + (n-1)d]$
 $\Rightarrow 2a_1(n-m) = d(n-m) \Rightarrow d = 2a_1$
 $\therefore \frac{a_m}{a_n} = \frac{\text{mth term}}{\text{nth term}} = \frac{a_1 + (m-1)d}{a_1 + (n-1)d}$
 $= \frac{a_1 + (m-1) \cdot 2a_1}{a_1 + (n-1) \cdot 2a_1} = \frac{-a_1 + 2a_1m}{-a_1 + 2a_1n}$
 $= \frac{a_1(2m-1)}{a_1(2n-1)} = \frac{2m-1}{2n-1}$.

12. Let a and d be the first term and common difference respectively of the A.P.

$$S_{10} = \frac{10}{2} [2a + 9d]$$

$$S_5 = \frac{5}{2} [2a + 4d]$$

Given, $S_{10} = 4S_5$

$$\Rightarrow 5(2a + 9d) = 4 \times \frac{5}{2} [2a + 4d]$$

$$\Rightarrow 10a + 45d = 20a + 40d$$

$$\Rightarrow 5d = 10a$$

$$\Rightarrow \frac{a}{d} = \frac{5}{10} = \frac{1}{2} \Rightarrow a : d = 1 : 2$$

13. The numbers between 107 and 253 divisible by 5 are

110, 115, 120, , 245, 250.

This is an A.P with first term (a) = 110 and common difference (d) = 5. Let the last term be the n th term.

$$\therefore T_n = a + (n-1)d$$

$$\Rightarrow 110 + (n-1) \times 5 = 250$$

$$\Rightarrow 5n = 250 - 105 = 145$$

$$\Rightarrow n = 29$$

$$\therefore \text{Required sum} = \frac{n}{2} (a + T_n) = \frac{29}{2} (110 + 250)$$

$$= \frac{29}{2} \times 360 = 5220$$

14. Let the two A.P.'s be $a, a + d, a + 2d, \dots$ and $A, A + D, A + 2D, \dots$

Given, $\frac{n/2 [2a + (n-1)d]}{n/2 [2A + (n-1)D]} = \frac{7n+1}{4n+27}$

$$\Rightarrow \frac{2a + (n-1)d}{2A + (n-1)D} = \frac{7n+1}{4n+27} \dots(i)$$

Now, we have to find the ratio $\frac{t_{11}}{T_{11}} = \frac{a + 10d}{A + 10D} = \frac{2a + 20d}{2A + 20D}$

Putting $n = 21$ in (i), we get

$$\frac{2a + 20d}{2A + 20D} = \frac{7 \times 21 + 1}{4 \times 21 + 27} = \frac{147 + 1}{84 + 27} = \frac{148}{111}$$

$$\therefore t_{11} : T_{11} = 148 : 111$$

15. Let a be the first term and d the common difference of the given A.P.

Then, $S_n = \frac{n}{2} [2a + (n-1)d]$

$$S_{2n} = \frac{2n}{2} [2a + (2n-1)d]$$

$$S_{3n} = \frac{3n}{2} [2a + (3n-1)d]$$

Given, $S_{2n} = 3S_n$

$$\Rightarrow \frac{2n}{2} [2a + (2n-1)d] = \frac{3n}{2} [2a + (n-1)d]$$

$$\Rightarrow 4a + 4nd - 2d = 6a + 3nd - 3d$$

$$\Rightarrow d + nd = 2a$$

$$\Rightarrow a = \frac{d(n+1)}{2} \dots(ii)$$

Now, $\frac{S_{3n}}{S_n} = \frac{\frac{3n}{2} [2a + (3n-1)d]}{\frac{n}{2} [2a + (n-1)d]}$
 $= \frac{3[(n+1)d + (3n-1)d]}{[(n+1)d + (n-1)d]} \quad (\text{From (i)})$
 $= \frac{3nd + 3d + 9nd - 3d}{nd + d + nd - d}$
 $= \frac{12nd}{2nd} = 6$

16. For the 1st A.P., 2, 5, 8, 11, ,

First term (a_1) = 2, common difference (d_1) = 3

\therefore The sum of this A.P. to $2n$ terms

$$= \frac{2n}{2} [2a_1 + (2n-1)d_1]$$

$$= n[4 + (2n-1)3]$$

$$= 4n + 6n^2 - 3n = 6n^2 + n = n(6n + 1)$$

For the second A.P., 57, 59, 61, 63, ,

First term (b_1) = 57, common difference (d_2) = 2

\therefore The sum of this A.P. to n term is

$$\text{Sum}_n = \frac{n}{2} [2b_1 + (n-1)d_2]$$

$$= \frac{n}{2} [114 + (n-1)2]$$

$$= 57n + n^2 - n = n^2 + 56n = n(n + 56)$$

$$\text{Given, } S_{2n} = \text{Sum}_n$$

$$\Rightarrow n(6n + 1) = n(n + 56)$$

$$\Rightarrow 6n + 1 = n + 56 \Rightarrow 5n = 55 \Rightarrow n = 11.$$

17. Let the first term of the given A.P. be a and the common difference be d .

$$\text{Then, } S_1 = \frac{n_1}{2} [2a + (n_1 - 1)d]$$

$$S_2 = \frac{n_2}{2} [2a + (n_2 - 1)d]$$

$$S_3 = \frac{n_3}{2} [2a + (n_3 - 1)d]$$

$$\begin{aligned} \therefore \frac{S_1}{n_1} (n_2 - n_3) + \frac{S_2}{n_2} (n_3 - n_1) + \frac{S_3}{n_3} (n_1 - n_2) \\ = \frac{n_1}{n_1} [2a + (n_1 - 1)d] (n_2 - n_3) + \frac{n_2}{n_2} [2a + (n_2 - 1)d] \\ (n_3 - n_1) + \frac{n_3}{n_3} [2a + (n_3 - 1)d] (n_1 - n_2) \\ = \frac{1}{2} [2a(n_2 - n_3) + n_1(n_2 - n_3) - d(n_2 - n_3)] \\ + \frac{1}{2} [2a(n_3 - n_1) + n_2(n_3 - n_1) - d(n_3 - n_1)] \\ + \frac{1}{2} [2a(n_1 - n_2) + n_3(n_1 - n_2) - d(n_1 - n_2)] \\ = \frac{1}{2} [2an_2 - 2an_3 + 2an_3 - 2an_1 + 2an_1 - 2an_2] + (n_1n_2 \\ - n_1n_3 + n_2n_3 - n_1n_2 + n_3n_1 - n_3n_2) \\ - d(n_2 - n_3 + n_3 - n_1 + n_1 - n_2)] \\ = 0. \end{aligned}$$

18. n th term of an A.P. is given by

$$T_n = \text{Sum of } n \text{ terms} - \text{Sum of } (n - 1) \text{ terms}$$

$$= S_n - S_{n-1}$$

$$= nP + \frac{n}{2} (n - 1)Q - \left[(n - 1)P + \frac{(n - 1)}{2} (n - 2)Q \right]$$

$$= nP - nP + P + \frac{Q}{2} [n^2 - n - n^2 + 3n - 2]$$

$$= P + \frac{Q}{2} (2n - 2)$$

$$= P + (n - 1)Q.$$

$$\begin{aligned} \therefore \text{Common difference } (d) &= T_n - T_{n-1} \\ &= P + (n - 1)Q - (P + (n - 2)Q) \\ &= Q. \end{aligned}$$

19. Let $a_1, a_2, a_3, \dots, a_n$ be the n arithmetic means between 3 and 17. Then,

$$3, a_1, a_2, a_3, \dots, a_n, 17 \text{ form an A.P.}$$

Let d be the common difference of this A.P. Then,

$$a_1 = 3 + d \quad \text{and} \quad a_n = 17 - d$$

$$\text{Given, } \frac{a_n}{a_1} = \frac{3}{1} \Rightarrow \frac{17 - d}{3 + d} = \frac{3}{1}$$

$$\Rightarrow 17 - d = 9 + 3d$$

$$\Rightarrow 4d = 8 \Rightarrow d = 2.$$

Also 17 is the $(n + 2)$ th term of the given A.P.

$$\therefore 17 = 3 + (n + 2 - 1)2$$

$$\Rightarrow 17 = 3 + (n + 1)2$$

$$\Rightarrow 14 = (n + 1)2 \Rightarrow n + 1 = 7 \Rightarrow n = 6.$$

20. Given, $a_1, a_2, a_3, \dots, a_n$ are in A.P. and $a_1 = 0$.

Let d be the common difference of the the A.P. So,

$$a_2 = a_1 + d = d$$

$$\therefore a_3 = a_1 + 2d = 2d, a_4 = 3d, a_5 = 4d, \dots, a_n = (n - 1)d$$

$$\begin{aligned} \therefore \left(\frac{a_3}{a_2} + \frac{a_4}{a_3} + \dots + \frac{a_n}{a_{n-1}} \right) - a_2 \left(\frac{1}{a_2} + \frac{1}{a_3} + \dots + \frac{1}{a_{n-2}} \right) \\ = \left[\frac{2d}{d} + \frac{3d}{2d} + \dots + \frac{(n-1)d}{(n-2)d} \right] \end{aligned}$$

$$- d \left[\frac{1}{d} + \frac{1}{2d} + \dots + \frac{1}{(n-3)d} \right]$$

$$= \left[2 + \frac{3}{2} + \dots + \frac{n-1}{n-2} \right] - \left[1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n-3} \right]$$

$$= \left\{ (1+1) + \left(1 + \frac{1}{2} \right) + \dots + \left(1 + \frac{1}{n-2} \right) \right\}$$

$$- \left\{ 1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n-3} \right\}$$

$$= 1 + 1 + 1 \dots \text{ to } (n - 2) \text{ term}$$

$$- \left(1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{(n-3)} + \frac{1}{(n-2)} \right)$$

$$- \left(1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n-3} \right)$$

$$= (n - 2) + \frac{1}{(n - 2)}.$$

21. Let $a_1 = a - 3d, a_2 = a - d, a_3 = a + d, a_4 = a + 3d$

$$\text{Given, } a_1 + a_4 = 10 \Rightarrow a - 3d + a + 3d = 10$$

$$\Rightarrow 2a = 10 \Rightarrow a = 5$$

$$\text{Also, } a_2 a_3 = 24 \Rightarrow (a - d)(a + d) = 24$$

$$\Rightarrow a^2 - d^2 = 24$$

$$\Rightarrow 25 - d^2 = 24 \Rightarrow d^2 = 1 \Rightarrow d = 1$$

$$\therefore \text{The least term is } a - 3d = 5 - 3 = 2.$$

22. Sum of first 10 terms = $\frac{1}{2}$ (Sum of next 10 terms)

$$\Rightarrow S_{10} = \frac{1}{2} (S_{20} - S_{10}) \Rightarrow 3S_{10} = S_{20}$$

$$\Rightarrow 3 \times \frac{10}{2} [2a + 9d] = \frac{20}{2} [2a + 19d]$$

$$\left[\because S_n = \frac{n}{2} [2a + (n-1)d] \right]$$

$$\Rightarrow 3(2a + 9d) = 2(2a + 19d)$$

$$\Rightarrow 6a + 27d = 4a + 38d$$

$$\Rightarrow 2a = 11d$$

$$\Rightarrow 2(13 + d) = 11d (\because \text{Second term} = 13 \text{ and } 13 - a = d)$$

$$\Rightarrow 26 - 2d = 11d$$

$$\Rightarrow 13d = 26 \Rightarrow d = 2.$$

23. Let the sides of the triangle be $a - d, a, a + d$.

Perimeter of the triangle = $a - d + a + a + d = 3a$

\therefore Each side of the equilateral triangle = $\frac{3a}{3} = a$

\therefore Area of equilateral triangle = $\frac{\sqrt{3}}{4} a^2$.

Area of the given $\Delta = \sqrt{s(s - (a - d))(s - a)(s - (a + d))}$

where $s = \frac{3a}{2}$

$$= \sqrt{\frac{3a}{2} \left(\frac{3a}{2} - a + d \right) \left(\frac{3a}{2} - a \right) \left(\frac{3a}{2} - a - d \right)}$$

$$= \sqrt{\frac{3a}{2} \left(\frac{1}{2} a + d \right) \left(\frac{1}{2} a \right) \left(\frac{1}{2} a - d \right)}$$

$$= \sqrt{\frac{3}{4} a^2 \left(\frac{1}{4} a^2 - d^2 \right)}$$

Given, $\sqrt{\frac{3}{4} a^2 \left(\frac{1}{4} a^2 - d^2 \right)} = \frac{3}{5} \times \frac{\sqrt{3}}{4} a^2$

$$\Rightarrow \frac{3}{16} a^4 - \frac{3}{4} a^2 d^2 = \left(\frac{3\sqrt{3}}{20} a^2 \right)^2$$

$$\Rightarrow \frac{3}{16} a^4 - \frac{3}{4} a^2 d^2 = \frac{27}{400} a^4$$

$$\Rightarrow \frac{3}{16} a^4 - \frac{27}{400} a^4 = \frac{3}{4} a^2 d^2$$

$$\Rightarrow \frac{1}{16} a^2 - \frac{9}{400} a^2 = \frac{1}{4} d^2$$

$$\Rightarrow \frac{25a^2 - 9a^2}{400} = \frac{1}{4} d^2$$

$$\Rightarrow \frac{16a^2}{400} = \frac{1}{4} d^2$$

$$\Rightarrow \frac{a^2}{d^2} = \frac{400}{64} \Rightarrow \frac{a}{d} = \frac{20}{8} = k \text{ (say)}$$

$$\therefore (a - d) : a : (a + d) = (20k - 8k) : 20k : (20k + 8k) = 12k : 20k : 28k = 3 : 5 : 7.$$

24. $\log_3 2, \log_3 (2^x - 5)$ and $\log_3 \left(2^x - \frac{7}{2} \right)$ are in A.P.

$$\Rightarrow 2 \log_3 (2^x - 5) = \log_3 2 + \log_3 (2^x - 7/2)$$

$$(\because a, b, c \text{ in A.P.} \Rightarrow 2b = a + c)$$

$$\Rightarrow \log_3 (2^x - 5)^2 = \log_3 [2(2^x - 7/2)]$$

$$(\because a \log b = \log b^a \text{ and } \log a + \log b = \log ab)$$

$$\Rightarrow (2^x - 5)^2 = 2^{x+1} - 7$$

Let $2^x = y$. Then,

$$(y - 5)^2 = 2y - 7$$

$$\Rightarrow y^2 - 10y + 25 = 2y - 7$$

$$\Rightarrow y^2 - 12y + 32 = 0$$

$$\Rightarrow (y - 8)(y - 4) = 0$$

$$\Rightarrow y = 8 \text{ or } 4 \Rightarrow 2^x = 8 \text{ or } 2^x = 4$$

$$2^x = 8 \Rightarrow x = 3$$

$$(\because 2^x = 4 \text{ shall make the term } \log_3(2^x - 5) \text{ negative which is not possible)}$$

25. $d = a_2 - a_1 = a_3 - a_2 = a_4 - a_3 = \dots = a_n - a_{n-1}$.

Now, $\sin d (\operatorname{cosec} a_1 \operatorname{cosec} a_2 + \operatorname{cosec} a_2 \operatorname{cosec} a_3 + \dots$

$+ \operatorname{cosec} a_{n-1} \operatorname{cosec} a_n)$

$$= \frac{\sin d}{\sin a_1 \sin a_2} + \frac{\sin d}{\sin a_2 \sin a_3} + \dots + \frac{\sin d}{\sin a_{n-1} \sin a_n}$$

$$= \frac{\sin (a_2 - a_1)}{\sin a_1 \sin a_2} + \frac{\sin (a_3 - a_2)}{\sin a_2 \sin a_3} + \dots + \frac{\sin (a_n - a_{n-1})}{\sin a_{n-1} \sin a_n}$$

$$= \frac{\sin a_2 \cos a_1 - \cos a_2 \sin a_1}{\sin a_1 \sin a_2}$$

$$+ \frac{\sin a_3 \cos a_2 - \cos a_3 \sin a_2}{\sin a_2 \sin a_3} + \dots$$

$$+ \frac{\sin a_n \cos a_{n-1} - \cos a_n \sin a_{n-1}}{\sin a_{n-1} \sin a_n}$$

$$= \frac{\cos a_1}{\sin a_1} - \frac{\cos a_2}{\sin a_2} + \frac{\cos a_2}{\sin a_2} - \frac{\cos a_3}{\sin a_3} + \dots$$

$$+ \frac{\cos a_{n-1}}{\sin a_{n-1}} - \frac{\cos a_n}{\sin a_n}$$

$$= \cot a_1 - \cot a_n.$$

GEOMETRIC PROGRESSION (G.P.)

KEY FACTS

1. A Geometric Progression (G.P.) is one in which the ratio of any term to its predecessor is always the same number. This ratio is called the **common ratio**. Thus, a sequence, a_1, a_2, \dots, a_n is said to be in G.P.,

$$\text{if } \frac{a_2}{a_1} = \frac{a_3}{a_2} = \frac{a_4}{a_3} = \dots = \frac{a_n}{a_{n-1}} = r$$

where a_1, a_2, a_3, \dots are all non-zero terms of the G.P. and r is the **common ratio**.

Examples of G.P.

Progression	Common ratio
(i) 1, 3, 9, 27, 81,	$r = 3$
(ii) $16/27, -8/9, 4/3, -2, \dots$	$r = -3/2$
(iii) x, x^2, x^3, x^4, \dots	$r = x$

If a denotes the first term of a G.P. and r , the common ratio, then a standard G.P. is a, ar, ar^2, \dots .

2. The n th term of a G.P.

For a standard G.P with **first term = a** , **common ratio = r** and **number of terms = n** , we know that:

First term	Second term	Third term	Fourth term	
$t_1 = a$	$t_2 = ar$	$t_3 = ar^2$	$t_4 = ar^3$	and so on.

So,

$$n\text{th term} = t_n = ar^{n-1}.$$

Ex. The sixth of the G.P. $-3, 6, -12, 24, \dots$ is:

Here $a = -3, r = \frac{6}{-3} = -2$

$$\therefore t_6 = ar^{6-1} = -3 \times (-2)^5 = -3 \times -32 = 96.$$

3. Geometric Mean: When three real numbers form a geometric progression, then the middle one is called the geometric mean of the other two quantities.

Thus, if G is the geometric mean of two non-zero numbers a and b , then a, G, b forms a geometric progression,

so that $\frac{G}{a} = \frac{b}{G}$

\therefore

$$G^2 = ab \Rightarrow G = \pm \sqrt{ab}$$

Maths Alert. $G = \sqrt{ab}$ if a and b are positive numbers.

$$G = -\sqrt{ab} \text{ if } a \text{ and } b \text{ are negative numbers.}$$

Also, if there are n positive integers a_1, a_2, \dots, a_n , then their geometric mean is defined to be equal to

$$(a_1 \cdot a_2 \cdot a_3 \cdot \dots \cdot a_n)^{1/n}.$$

Ex. (i) G.M. of 4 and 36 = $\sqrt{4 \times 36} = \sqrt{144} = 12$.

$$\begin{aligned} \text{(ii) G.M. of } \frac{1}{3}, 1, 3, 9, 27, 81 \text{ and } 243 &= \left(\frac{1}{3} \times 1 \times 3 \times 9 \times 27 \times 81 \times 243 \right)^{1/7} \\ &= (3^2 \times 3^3 \times 3^4 \times 3^5)^{1/7} = (3^{14})^{1/7} = 3^2 = 9. \end{aligned}$$

4. Sum of n terms of a Geometric Progression:

Consider the geometric progression $a, ar, ar^2, \dots, ar^{n-1}$ of n terms.

Then, the sum of n terms $S_n = \frac{a(1-r^n)}{1-r} (r \neq 1)$ or $\frac{a(r^n-1)}{(r-1)} (r \neq 1)$

where, a = first term, r = common ratio, n = number of terms

Also, $S_n = \frac{a - lr}{1 - r}$ or $\frac{lr - a}{r - 1}$, where l is the last term.

Ex. The sum of the G.P. $4 + 2 + 1 + \frac{1}{2} + \frac{1}{4} + \dots$ to 10 term is:

Here $a = 4, r = \frac{1}{2}, n = 10$

$$\therefore S_{10} = \frac{4 \left[1 - \left(\frac{1}{2} \right)^{10} \right]}{1 - \frac{1}{2}} = \frac{4 \left(1 - \frac{1}{1024} \right)}{\frac{1}{2}} = \frac{8 \times 1023}{1024} = 8 \text{ (approx.)}$$

5. Sum of an infinite G.P.

$S_\infty = \frac{a}{1 - r}$, where a = first term
 r = common ratio.

Note: Sum of an infinite series exists only when r is numerically less than 1, i.e., $|r| < 1$.

Ex. Sum of the G.P. $16, -8, 4, \dots$ to infinity is:

Here $a = 16, r = -\frac{8}{16} = -\frac{1}{2}$

$$\therefore S_\infty = \frac{a}{1 - r} = \frac{16}{1 - \left(-\frac{1}{2} \right)} = \frac{16}{1 + \frac{1}{2}} = \frac{16}{\frac{3}{2}} = \frac{32}{3}$$

6. Properties of a G.P.

(a) If each term of a given geometric progression is multiplied or divided by the same number, then the resulting progression is also a G.P.

If $a_1, a_2, a_3, a_4, \dots$ be a G.P. with common ratio r , then

$a_1c, a_2c, a_3c, a_4c, \dots$ and $\frac{a_1}{c}, \frac{a_2}{c}, \frac{a_3}{c}, \frac{a_4}{c}, \dots$ are also G.P.s with common ratio r . c is the constant number.

(b) The reciprocals of the terms of a G.P. also form a G.P.

If $a_1, a_2, a_3, a_4, \dots$ be a G.P. with common ratio r , then $\frac{1}{a_1}, \frac{1}{a_2}, \frac{1}{a_3}, \frac{1}{a_4}, \dots$ is also a G.P. with common ratio $\frac{1}{r}$.

(c) If each term of a G.P. be raised to the same power, then the resulting progression is also a G.P.

If $a_1, a_2, a_3, a_4, \dots$ be a G.P. with common ratio r , then $a_1^k, a_2^k, a_3^k, a_4^k, \dots$ is also a G.P. with common ratio r .

(d) If $a_1, a_2, a_3, a_4, \dots$ and $b_1, b_2, b_3, b_4, \dots$ be two G.P.s with common ratio r_1 and r_2 respectively, then the

progressions $a_1b_1, a_2b_2, a_3b_3, \dots$ is a G.P. with common ratio r_1r_2 , and $\frac{a_1}{b_1}, \frac{a_2}{b_2}, \frac{a_3}{b_3}, \frac{a_4}{b_4}, \dots$ is a G.P. with

common ratio $\frac{r_1}{r_2}$.

(e) If a_1, a_2, a_3, \dots be a G.P. with common ratio r , such that each term of the progression is a positive number, then $\log a_1, \log a_2, \log a_3, \dots$ is an A.P. with common difference $\log r$.

Conversely, if $\log a_1, \log a_2, \log a_3, \dots$ are terms of an A.P., then a_1, a_2, a_3, \dots are terms of a G.P.

(f) The Arithmetic mean between two positive numbers is always greater than equal to their Geometric Mean.

$$A.M. \geq G.M.$$

(g) If A and G be the arithmetic and geometric mean respectively between two numbers, then the numbers are

$$A \pm \sqrt{A^2 - G^2}.$$

7. Useful note: When product is known, suppose the numbers in G.P. as under:

(i) If the odd number of terms are to be considered, suppose them as follows by taking a as the mid-term and common ratio r .

$$\frac{a}{r}, a, ar; \quad \frac{a}{r^2}, \frac{a}{r}, a, ar, ar^2$$

(ii) If even number of terms are to be considered, suppose them as under by taking $\frac{a}{r}$, ar as the two mid-terms and r^2 as the common ratio.

$$\frac{a}{r^3}, \frac{a}{r}, ar, ar^3 \quad [4 \text{ terms}]$$

$$\frac{a}{r^5}, \frac{a}{r^3}, \frac{a}{r}, ar, ar^3, ar^5 \quad [6 \text{ terms}]$$

[See Solved Ex. (7)]

SOLVED EXAMPLES

Ex. 1. Write the G.P. whose 4th term is 54 and 7th term is 1458.

Sol. $t_4 = ar^3 = 54$...(i)

$t_7 = ar^6 = 1458$...(ii)

\therefore Dividing (ii) by (i), we get

$$\frac{t_7}{t_4} = r^3 = \frac{1458}{54} = 27$$

$\Rightarrow r = 3$

\therefore From (i), $a(3)^3 = 54$

$\Rightarrow 27a = 54$

$\Rightarrow a = 2$

\therefore The G.P. is $2, 2 \times 3, 2 \times 3^2, \dots$, i.e. $2, 6, 18, 54, \dots$.

Ex. 2. If the 10th term of a G.P. is 9 and the 4th term is 4, then what is its 7th term?

Sol. Let the first term and common ratio of the given G.P. be a and r respectively.

$T_{10} = ar^9 = 9$...(i)

$T_4 = ar^3 = 4$...(ii)

$$\frac{T_{10}}{T_4} = \frac{ar^9}{ar^3} = \frac{9}{4}$$

$r^6 = \frac{9}{4} \Rightarrow (r^3)^2 = \frac{9}{4} \Rightarrow r^3 = \frac{3}{2}$...(iii)

From (i), $ar^9 = 9 \Rightarrow a(r^3)^3 = 9$

\therefore Substituting the value of r^3 from (iii), we get

$$a \cdot \left(\frac{3}{2}\right)^3 = 9 \Rightarrow a = \frac{9 \times 8}{27} = \frac{8}{3}$$

$\therefore T_7 = ar^6 = \frac{8}{3} \times \frac{9}{4} = 6.$

Ex. 3. What is the sum of 10 terms of the series $\sqrt{2} + \sqrt{6} + \sqrt{18} + \dots$?

(DCE)

Sol.
$$S_n = \frac{a(r^n - 1)}{(r - 1)}$$

Here $a = \sqrt{2}, r = \frac{\sqrt{6}}{\sqrt{2}} = \sqrt{3}, n = 10$

$$\begin{aligned} \therefore S_{10} &= \frac{\sqrt{2}((\sqrt{3})^{10} - 1)}{\sqrt{3} - 1} = \frac{\sqrt{2}(3^5 - 1)}{\sqrt{3} - 1} \times \frac{\sqrt{3} + 1}{\sqrt{3} + 1} = \frac{\sqrt{2}(243 - 1)(\sqrt{3} + 1)}{(3 - 1)} \\ &= \frac{\sqrt{2}(242)(\sqrt{3} + 1)}{2} = 121(\sqrt{6} + \sqrt{2}). \end{aligned}$$

Ex. 4. What is the sum to infinity of the series $1 - \frac{1}{2} + \frac{1}{4} - \frac{1}{8} + \dots$?

(NDA/NA 2012)

Sol. Given series is an infinite G.P. with $a = 1, r = -\frac{1}{2}/1 = -\frac{1}{2}$.

$$\begin{aligned} \therefore S_{\infty} &= \frac{a}{1 - r} \\ &= \frac{1}{1 - \left(-\frac{1}{2}\right)} = \frac{1}{\frac{3}{2}} = \frac{2}{3}. \end{aligned}$$

Ex. 5. If a, b, c are the three consecutive terms of an A.P. and x, y, z are three consecutive terms of a G.P., then prove that $x^{b-c} \cdot y^{c-a} \cdot z^{a-b} = 1$.

Sol. a, b, c are in A.P. $\Rightarrow 2b = a + c$...(i)

x, y, z are in G.P. $\Rightarrow y = xr, z = xr^2$, where r is the common ratio

$$\begin{aligned} \therefore x^{b-c} \cdot y^{c-a} \cdot z^{a-b} &= x^{b-c} \cdot (xr)^{c-a} \cdot (xr^2)^{a-b} \\ &= x^{b-c} \cdot x^{c-a} \cdot r^{c-a} \cdot x^{a-b} \cdot r^{2a-2b} = x^{b-c+c-a+a-b} \cdot r^{c-a+2a-2b} \\ &= x^0 \cdot r^{c+a-2b} = x^0 \cdot r^{2b-2b} = x^0 \cdot r^0 = 1 \end{aligned} \quad \text{(From (i))}$$

Ex. 6. Let two numbers have arithmetic mean 9 and geometric mean 4. Then find the equation which has these two numbers as its roots. (AIEEE 2004)

Sol. Let the two numbers be a and b .

Then $\frac{a+b}{2} = 9 \Rightarrow a+b = 18$

$$\sqrt{ab} = 4 \Rightarrow ab = 16$$

$$\begin{aligned} \therefore \text{Required eqn is } x^2 - \text{Sum of roots } x + \text{Product of roots} &= 0 \\ \Rightarrow x^2 - 18x + 16 &= 0. \end{aligned}$$

Ex. 7. Insert 3 geometric means between 16 and 256.

Sol. Let G_1, G_2, G_3 be the required means.

Then 16, $G_1, G_2, G_3, 256$ form a G.P.

Let r be the common ratio.

$$\Rightarrow 256 = 5\text{th term} = ar^4 = 16 \times r^4$$

$$\Rightarrow 16r^4 = 256 \Rightarrow r^4 = 16 \Rightarrow r = 2$$

$$\therefore G_1 = ar = 16 \times 2 = 32$$

$$G_2 = ar^2 = 16 \times 4 = 64$$

$$G_3 = ar^3 = 16 \times 8 = 128$$

Hence 32, 64 and 128 are the required G.M's between 16 and 256.

Ex. 8. If one AM 'A' and two GM p and q are inserted between two given numbers, then find the value of $\frac{p^2}{q} + \frac{q^2}{p}$ in terms of A ? (VITEEE 2011)

Sol. Let a and b be the two given numbers.

$$\text{Then, A.M.} = \frac{a+b}{2} \Rightarrow A = \frac{a+b}{2} \Rightarrow 2A = a+b \quad \dots(i)$$

As, p and q are G.M's between a and b , a, p, q, b forms a G.P.

$$\therefore \frac{p}{a} = \frac{q}{p} = \frac{b}{q} \Rightarrow \frac{p^2}{a} = b \text{ and } \frac{q^2}{b} = a$$

$$\text{Now, } \frac{p^2}{q} + \frac{q^2}{p} = a + b = 2A.$$

Ex. 9. What is the sum of the series $\frac{2}{3} + \frac{8}{9} + \frac{26}{27} + \frac{80}{81} + \dots$ to n terms? (KCET 2001)

Sol. The series $\frac{2}{3} + \frac{8}{9} + \frac{26}{27} + \frac{80}{81} + \dots$ to n terms can be written as

$$\begin{aligned} & \left(1 - \frac{1}{3}\right) + \left(1 - \frac{1}{9}\right) + \left(1 - \frac{1}{27}\right) + \left(1 - \frac{1}{81}\right) + \dots \text{ to } n \text{ terms} \\ &= (1 + 1 + 1 + 1 + \dots \text{ to } n \text{ terms}) - \left(\frac{1}{3} + \frac{1}{9} + \frac{1}{27} + \frac{1}{81} + \dots \text{ to } n \text{ terms}\right) \\ &= n - \frac{1 \left(1 - \left(\frac{1}{3}\right)^n\right)}{1 - \frac{1}{3}} \quad \left(S_n = \frac{a(1-r^n)}{1-r}, \text{ here } a = \frac{1}{3}, r = \frac{1}{3}\right) \\ &= n - \frac{1 \left(1 - \frac{1}{3^n}\right)}{2/3} = n - \frac{1}{2}(1 - 3^{-n}). \end{aligned}$$

Ex. 10. If $1 + \sin x + \sin^2 x + \sin^3 x + \dots \infty = 4 + 2\sqrt{3}$, $0 < x < \pi$, then find the value of x . (Kerala PET 2004)

Sol. Given, $1 + \sin x + \sin^2 x + \sin^3 x + \dots \infty = 4 + 2\sqrt{3}$

$$\Rightarrow \frac{1}{1 - \sin x} = 4 + 2\sqrt{3}$$

$$\Rightarrow 1 - \sin x = \frac{1}{4 + 2\sqrt{3}}$$

$$\begin{aligned} \Rightarrow \sin x &= 1 - \frac{1}{4 + 2\sqrt{3}} = \frac{4 + 2\sqrt{3} - 1}{4 + 2\sqrt{3}} = \frac{3 + 2\sqrt{3}}{4 + 2\sqrt{3}} \times \frac{4 - 2\sqrt{3}}{4 - 2\sqrt{3}} \\ &= \frac{12 + 8\sqrt{3} - 6\sqrt{3} - 12}{16 - 12} = \frac{2\sqrt{3}}{4} = \frac{\sqrt{3}}{2} \end{aligned}$$

$$\Rightarrow x = \frac{\pi}{3} \text{ or } \frac{2\pi}{3} \text{ as } 0 < x < \pi.$$

$$\left[\begin{array}{l} \because S_\infty = \frac{a}{1-r}. \text{ Here } a = 1, r = \sin x \\ \text{Also } |\sin x| < 1 \text{ as } |\sin x| = 1 \text{ will make} \\ \text{the the sum of infinte G.P. infinite) } \end{array} \right]$$

Ex. 11. If a, b, c are in G.P. and $4a, 5b, 4c$ are in A.P. such that $a + b + c = 70$, then what is the value of the smallest of the numbers a, b and c ?

Sol. a, b, c are in G.P. $\Rightarrow b^2 = ac$...(i)

$4a, 5b, 4c$ are in A.P. $\Rightarrow 2 \times 5b = 4a + 4c \Rightarrow 10b = 4a + 4c \Rightarrow 5b = 2a + 2c$...(ii)

Also, given $a + b + c = 70$...(iii)

$\Rightarrow 2a + 2b + 2c = 140 \Rightarrow 5b + 2b = 140$ (From (ii))

$\Rightarrow 7b = 140 \Rightarrow b = 20.$

Now, from (i), $400 = ac$. ($\because b = 20$)

Also, from (ii), $a + 20 + c = 70 \Rightarrow a + c = 50$

$\therefore (a - c)^2 = (a + c)^2 - 4ac$
 $= 2500 - 1600 = 900$

$\Rightarrow a - c = \pm 30$

$\therefore \left. \begin{array}{l} a + c = 50 \\ a - c = \pm 30 \end{array} \right\} \Rightarrow a = 40, c = 10 \text{ or } a = 10, c = 40.$

\therefore The least value out of a, b and c is **10**.

Ex. 12. Find the sum of n terms of the series $1 + (1 + x) + (1 + x + x^2) + \dots$?

(AMU 2003)

Sol. $1 + (1 + x) + (1 + x + x^2) + \dots$ n terms

\Rightarrow Required sum $= \frac{1}{(1-x)} [(1-x) + (1-x)(1+x) + (1-x)(1+x+x^2) + \dots n \text{ terms}]$

$= \frac{1}{(1-x)} [(1-x) + (1-x^2) + (1-x^3) + \dots n \text{ terms}]$

$= \frac{1}{(1-x)} [(1 + 1 + 1 + \dots n \text{ terms}) - (x + x^2 + x^3 + \dots n \text{ terms})]$

$= \frac{1}{(1-x)} \left[n - \frac{x(1-x^n)}{(1-x)} \right]$ ($\because S_n = \frac{a(1-r^n)}{(1-r)}$, Here $a = x, r = x$)

$= \frac{n(1-x) - x(1-x^n)}{(1-x)^2}.$

Ex. 13. Five numbers are in A.P. with common difference $\neq 0$. If 1st, 3rd and 4th terms are in G.P., then which term is always zero? (WBJEE 2013)

Sol. Let the five numbers in A.P. be $a - 2d, a - d, a, a + d, a + 2d$.

Since 1st, 3rd and 4th terms are in G.P.

$(t_3)^2 = t_1 \cdot t_4$

$\Rightarrow a^2 = (a - 2d)(a + d)$

$\Rightarrow a^2 = a^2 - ad - 2d^2$

$\Rightarrow -ad = 2d^2$

$\Rightarrow a = -2d$ ($\because d \neq 0$)

$\Rightarrow a + 2d = 0 \Rightarrow t_5 = 0$

\therefore 5th term is always zero.

Ex. 14. Find the sum to n terms of the series, $7 + 77 + 777 + \dots$

Sol. $S_n = 7 + 77 + 777 + \dots$ to n terms
 $= 7 [1 + 11 + 111 + \dots$ to n terms]
 $= \frac{7}{9} [9 + 99 + 999 + \dots$ to n terms]

$$\begin{aligned}
&= \frac{7}{9} [(10 - 1) + (100 - 1) + (1000 - 1) + \dots \text{ to } n \text{ terms}] \\
&= \frac{7}{9} [\{10 + 10^2 + 10^3 + \dots \text{ to } n \text{ terms}\} - \{1 + 1 + 1 + \dots \text{ to } n \text{ terms}\}] \\
&= \frac{7}{9} \left[\frac{10(10^n - 1)}{10 - 1} - n \right] = \frac{7(10^{n+1} - 10)}{81} - \frac{7}{9}n.
\end{aligned}$$

Ex. 15. What is the sum of n terms of the series $0.2 + 0.22 + 0.222 + \dots$?

(WBJEE 2009)

Sol. $0.2 + 0.22 + 0.222 + \dots$ to n terms

$$\begin{aligned}
&= 2[0.1 + 0.11 + 0.111 + \dots \text{ to } n \text{ terms}] \\
&= \frac{2}{9} [0.9 + 0.99 + 0.999 + \dots \text{ to } n \text{ terms}] \\
&= \frac{2}{9} [(1 - 0.1) + (1 - 0.01) + (1 + 0.001) + \dots \text{ to } n \text{ terms}] \\
&= \frac{2}{9} [(1 + 1 + 1 + \dots \text{ to } n \text{ terms}) - (0.1 + 0.01 + 0.001 + \dots \text{ to } n \text{ terms})] \\
&= \frac{2}{9} \left[n - \frac{0.1(1 - (0.1)^n)}{(1 - 0.1)} \right] = \frac{2}{9} \left[n - \frac{\frac{1}{10} \left(1 - \frac{1}{10^n} \right)}{1 - \frac{1}{10}} \right] = \frac{2}{9} \left[n - \frac{1}{9} \left(1 - \frac{1}{10^n} \right) \right].
\end{aligned}$$

Ex. 16. Find the value of $0.2\overline{34}$ regarding it as a geometric series.

Sol.

$$\begin{aligned}
0.2\overline{34} &= 0.234\ 34\ 34\ \dots \\
&= 0.2 + 0.034 + 0.00034 + 0.0000034 + \dots + \infty \\
&= \frac{2}{10} + \frac{34}{1000} + \frac{34}{100000} + \frac{34}{10000000} + \dots + \infty \\
&= \frac{2}{10} + \frac{34}{10^3} \left[1 + \frac{1}{10^2} + \frac{1}{10^4} + \dots \infty \right] \\
&= \frac{2}{10} + \frac{34}{10^3} \times \left(\frac{1}{1 - \frac{1}{10^2}} \right) \quad \left(\because S_{\infty} = \frac{a}{1 - r}. \text{ Here } a = 1, r = \frac{1}{10^2} \right) \\
&= \frac{2}{10} + \frac{34}{1000} \times \frac{100}{99} = \frac{198 + 34}{990} = \frac{232}{990} = \frac{116}{495}.
\end{aligned}$$

Ex. 17. Six positive numbers are in G.P. such that their product is 1000. If the fourth term is 1, then find the last term. (WBJEE 2013)

Sol. Let the six numbers in G.P. be $\frac{a}{r^5}, \frac{a}{r^3}, \frac{a}{r}, ar, ar^3, ar^5$.

Then, Product = 1000

$$\Rightarrow \frac{a}{r^5} \times \frac{a}{r^3} \times \frac{a}{r} \times ar \times ar^3 \times ar^5 = 1000$$

$$\Rightarrow a^6 = 1000 \Rightarrow a = \sqrt[6]{1000}$$

Given, Fourth term = $t_4 = ar = 1$

$$\Rightarrow \sqrt[6]{1000} \times r = 1 \Rightarrow r = \frac{1}{\sqrt[6]{1000}}$$

$$\therefore \text{Last term} = ar^5 = \sqrt[6]{1000} \times \left(\frac{1}{\sqrt[6]{1000}} \right)^5 = \frac{1}{100}.$$

Ex. 18. If a, b, c, d are in G.P., prove that $(a^2 + b^2 + c^2), (ab + bc + cd), (b^2 + c^2 + d^2)$ are in G.P.

Sol. a, b, c, d are in G.P $\Rightarrow b = ar, c = ar^2, d = ar^3$, where $r =$ common ratio

$$\therefore (ab + bc + cd)^2 = (a \cdot ar + ar \cdot ar^2 + ar^2 \cdot ar^3)^2 = [a^2r(1 + r^2 + r^4)]^2 \quad \dots(i)$$

$$\begin{aligned} (a^2 + b^2 + c^2)(b^2 + c^2 + d^2) &= (a^2 + a^2r^2 + a^2r^4)(a^2r^2 + a^2r^4 + a^2r^6) \\ &= a^2(1 + r^2 + r^4) \cdot a^2r^2(1 + r^2 + r^4) = a^4r^2(1 + r^2 + r^4)^2 \\ &= [a^2r(1 + r^2 + r^4)]^2 = (ab + bc + cd)^2 \end{aligned}$$

(From (i))

$\therefore (a^2 + b^2 + c^2), (ab + bc + cd), (b^2 + c^2 + d^2)$ are in G.P.

Ex. 19. If x, y, z are all positive and are the p th, q th and r th terms of a geometric progression respectively, then

find the value of the determinant $\begin{vmatrix} \log x & p & 1 \\ \log y & q & 1 \\ \log z & r & 1 \end{vmatrix}$. (VITEEE 2009)

Sol. Let a be the first term and R the common ratio of the G.P.

Then,

$$T_p = aR^{p-1} = x$$

$$T_q = aR^{q-1} = y$$

$$T_r = aR^{r-1} = z$$

$$\Rightarrow \log x = \log(aR^{p-1}) = \log a + (p-1) \log R$$

$$\log y = \log(aR^{q-1}) = \log a + (q-1) \log R$$

$$\log z = \log(aR^{r-1}) = \log a + (r-1) \log R$$

$$\therefore \begin{vmatrix} \log x & p & 1 \\ \log y & q & 1 \\ \log z & r & 1 \end{vmatrix} = \begin{vmatrix} \log a + (p-1) \log R & p & 1 \\ \log a + (q-1) \log R & q & 1 \\ \log a + (r-1) \log R & r & 1 \end{vmatrix}$$

$$= \begin{vmatrix} \log a & p & 1 \\ \log a & q & 1 \\ \log a & r & 1 \end{vmatrix} + \begin{vmatrix} (p-1) \log R & p & 1 \\ (q-1) \log R & q & 1 \\ (r-1) \log R & r & 1 \end{vmatrix}$$

$$= \log a \begin{vmatrix} 1 & p & 1 \\ 1 & q & 1 \\ 1 & r & 1 \end{vmatrix} + \log R \begin{vmatrix} (p-1) & p & 1 \\ (q-1) & q & 1 \\ (r-1) & r & 1 \end{vmatrix}$$

$$= \log a \times 0 + \log R \begin{vmatrix} p & p & 1 \\ q & q & 1 \\ r & r & 1 \end{vmatrix} - \log R \begin{vmatrix} 1 & p & 1 \\ 1 & q & 1 \\ 1 & r & 1 \end{vmatrix}$$

$$= 0 + \log R \times 0 - \log R \times 0 = 0$$

(\because The value of a determinants with two identical rows or columns is zero).

Ex. 20. If $A = 1 + r^a + r^{2a} + r^{3a} + \dots \infty, B = 1 + r^b + r^{2b} + r^{3b} + \dots \infty$, then show that $\frac{a}{b} = \log \left(\frac{B-1}{B} \right) \left(\frac{A-1}{A} \right)$.

Sol. $A = 1 + r^a + r^{2a} + r^{3a} + \dots \infty$

$$\Rightarrow A = \frac{1}{1-r^a} \quad \left(\because S_\infty = \frac{a}{1-r}. \text{ Here } a=1, \text{ common ratio } = r^a \right)$$

$$\Rightarrow 1 - r^a = \frac{1}{A} \Rightarrow r^a = 1 - \frac{1}{A} = \frac{A-1}{A} \Rightarrow a \log r = \log \left(\frac{A-1}{A} \right)$$

$$\Rightarrow a = \frac{\log \left(\frac{A-1}{A} \right)}{\log r} \quad \dots(i)$$

Now, $B = 1 + r^b + r^{2b} + r^{3b} + \dots \infty$

$$\Rightarrow B = \frac{1}{1-r^b} \Rightarrow 1-r^b = \frac{1}{B} \Rightarrow r^b = 1 - \frac{1}{B} = \frac{B-1}{B} \Rightarrow \log r^b = \log \left(\frac{B-1}{B} \right)$$

$$\Rightarrow b \log r = \log \left(\frac{B-1}{B} \right) \Rightarrow b = \frac{\log \left(\frac{B-1}{B} \right)}{\log r} \quad \dots(ii)$$

$$\therefore \frac{a}{b} = \frac{\log \left(\frac{A-1}{A} \right)}{\log r} \times \frac{\log r}{\log \left(\frac{B-1}{B} \right)} \quad \text{(From (i) and (ii))}$$

$$\Rightarrow \frac{a}{b} = \frac{\log \left(\frac{A-1}{A} \right)}{\log \left(\frac{B-1}{B} \right)} = \log_{\left(\frac{B-1}{B} \right)} \left(\frac{A-1}{A} \right). \quad \left[\because \frac{\log a}{\log b} = \log_b a \right]$$

Ex. 21. $S_1, S_2, S_3, \dots, S_n$ are the sums of n infinite geometric progressions. The first term S of these progressions are $1, 2^2 - 1, 2^3 - 1, 2^4 - 1, \dots, 2^n - 1$ and the common ratios are $\frac{1}{2}, \frac{1}{2^2}, \frac{1}{2^3}, \dots, \frac{1}{2^n}$. Calculate the sum $S_1 + S_2 + S_3 + \dots + S_n$.

Sol. Since $S_\infty = \frac{a}{1-r}$,

$$S_1 = \frac{1}{1 - \frac{1}{2}} = \frac{1}{\frac{1}{2}} = 2$$

$$S_2 = \frac{2^2 - 1}{1 - \frac{1}{2^2}} = \frac{2^2 - 1}{\frac{2^2 - 1}{2^2}} = 2^2$$

$$S_3 = \frac{2^3 - 1}{1 - \frac{1}{2^3}} = \frac{2^3 - 1}{\frac{2^3 - 1}{2^3}} = 2^3$$

.....

$$S_n = \frac{2^n - 1}{1 - \frac{1}{2^n}} = \frac{2^n - 1}{\frac{2^n - 1}{2^n}} = 2^n$$

$$\therefore S_1 + S_2 + S_3 + \dots + S_n = 2 + 2^2 + 2^3 + \dots + 2^n$$

$$= \frac{2(2^n - 1)}{2 - 1} = 2(2^n - 1)$$

$$\left(\because S_n = \frac{a(r^n - 1)}{r - 1}, \text{ Here } a = 2, r = 2 \right)$$

PRACTICE SHEET

Level-1

1. Which term of the G.P. $3, 3\sqrt{3}, 9, \dots$ is 2187?

- (a) 13 (b) 14 (c) 15 (d) 16

(Kerala 2004)

2. If $n!, 3 \times (n!)$ and $(n+1)!$ are in G.P, then the value of n will be

- (a) 3 (b) 4 (c) 8 (d) 10

(NDA/NA 2011)

3. If 1, $x, y, z, 16$ are in G.P., then what is the value of $x + y + z$?

- (a) 8 (b) 12 (c) 14 (d) 16

(NDA/NA 2008)

4. The geometric mean of $1, 2, 2^2, \dots, 2^n$ is

- (a) $2^{n/2}$ (b) $n^{(n+1)/2}$ (c) $2^{(n-1)/2}$ (d) $2^{n(n+1)/2}$

(MPPET 2009)

5. If the first term of a G.P. is 729 and its 7th term is 64, then the sum of the first seven terms is

- (a) 2187 (b) 2059 (c) 1458 (d) 2123

(Kerala 2013)

6. If the third term of a G.P. is 3, then the product of its first 5 terms is:

- (a) 15 (b) 81
(c) 243 (d) Cannot be determined

(J & K CET 2009)

7. In a G.P, $t_2 + t_5 = 216$ and $t_4 : t_6 = 1 : 4$ and all the terms are integers, then its first term is

- (a) 16 (b) 14 (c) 12 (d) None of these

(AMU 2010)

8. The sum of the first n terms of the series

$$\frac{1}{2} + \frac{3}{4} + \frac{7}{8} + \frac{15}{16} + \dots \text{ is}$$

- (a) $2^n - n - 1$ (b) $1 - 2^{-n}$
(c) $n + 2^{-n} - 1$ (d) $2^n - 1$ (AMU 2003)

9. In a geometric progression consisting of positive terms, each term equals the sum of next two terms. Then, the common ratio of the progression equals

- (a) $\frac{\sqrt{5}}{2}$ (b) $\sqrt{5}$ (c) $\frac{\sqrt{5}-1}{2}$ (d) $\frac{\sqrt{5}+1}{2}$

(AIEEE 2007)

10. If in an infinite G.P, the first term equal to twice the sum of all the successive terms, then the common of this G.P. is

- (a) $\frac{1}{3}$ (b) $\frac{1}{4}$ (c) $\frac{2}{5}$ (d) $\frac{2}{3}$

(Rajasthan 2002)

Level-2

11. Consider an infinite geometric series with first term a and common ratio r . If its sum is 4 and the second term is $\frac{3}{4}$, then

- (a) $a = 2, r = \frac{1}{2}$ (b) $a = 2, r = \frac{3}{8}$
(c) $a = 1, r = \frac{3}{4}$ (d) None of these

(AMU 2013)

12. If $x, 2x + 2, 3x + 3$, are the first three terms of a G.P, then the fourth term is

- (a) $\frac{-27}{2}$ (b) $\frac{27}{2}$ (c) $\frac{-33}{2}$ (d) $\frac{33}{2}$

(NDA/NA 2009, Rajasthan PET 2005)

13. Six positive numbers are in G.P, such that their product is 1000. If the fourth term is 1, then the last term is

- (a) 1000 (b) 100 (c) $\frac{1}{100}$ (d) $\frac{1}{1000}$

(WBJEE 2013)

14. If a is the A.M. of b and c and the two geometric means are G_1 and G_2 , then $G_1^3 + G_2^3$ is equal to

- (a) $\frac{abc}{2}$ (b) abc (c) $2abc$ (d) $\frac{3}{2}abc$

(BCECE 2009, IIT 1997)

15. If a, b, c are unequal numbers such that a, b, c are in A.P. and $b - a, c - b, a$ are in G.P, then $a : b : c$ is

- (a) $1 : 2 : 3$ (b) $1 : 2 : 4$ (c) $1 : 3 : 4$ (d) $2 : 3 : 4$

(AMU 2005)

16. What is the sum of the 100 terms of the series $9 + 99 + 999 + \dots$?

- (a) $\frac{10}{9}(10^{100} - 1) - 100$ (b) $\frac{10}{9}(10^{99} - 1) - 100$

- (c) $100(100^{10} - 1)$ (d) $\frac{9}{100}(10^{100} - 1)$

(NDA/NA 2008)

17. The sum of the first 20 terms of the sequence $0.7, 0.77, 0.777, \dots$ is

- (a) $\frac{7}{81}(179 - 10^{-20})$ (b) $\frac{7}{9}(99 - 10^{-20})$

- (c) $\frac{7}{81}(179 + 10^{-20})$ (d) $\frac{7}{9}(99 + 10^{-20})$

(IIT JEE 2013)

18. If $\frac{2}{3} = \left(x - \frac{1}{y}\right) + \left(x^2 - \frac{1}{y^2}\right) + \dots$ to ∞ and $xy = 2$, then the value of x and y under the condition $x < 1$ are

- (a) $x = \frac{1}{3}, y = 6$ (b) $x = \frac{1}{2}, y = 4$

- (c) $x = \frac{1}{4}, y = 8$ (d) $x = \frac{1}{6}, y = 12$

19. The first two terms of a geometric progression add upto 12. The sum of the third and fourth terms is 48. If the terms of the geometric progression are alternately positive and negative, then the first term is

- (a) -4 (b) -12 (c) 12 (d) 4

(AIEEE 2008)

20. If 64, 27 and 36 are the P th, Q th and R th terms of a G.P, then $P + 2Q$ is equal to

- (a) R (b) $2R$ (c) $3R$ (d) $4R$

(WBJEE 2012)

Level-3

21. In a sequence of 21 terms, the first 11 terms are in A.P. with common difference 2 and the last 11 terms are in G.P with common ratio 2. If the middle term of A.P. be equal to the middle term of G.P., then the middle term of the entire sequence is

- (a) $\frac{-10}{31}$ (b) $\frac{10}{31}$ (c) $\frac{32}{31}$ (d) $\frac{-31}{32}$

(AMU 2009)

22. If $x = 1 + a + a^2 + \dots \infty$ and $y = 1 + b + b^2 + \dots \infty$, where a and b are proper fractions, then $1 + ab + a^2b^2 + \dots \infty$ equals

(a) $\frac{x+y}{x-y}$ (b) $\frac{x^2+y^2}{x-y}$ (c) $\frac{x^2-y^2}{x+y-1}$ (d) $\frac{xy}{x+y-1}$

(Punjab CET 2007)

23. The first term of an infinite G.P. is x and its sum is 5. Then,

(a) $-10 < x < 0$ (b) $0 < x < 10$
(c) $0 \leq x \leq 10$ (d) $x > 10$

(IIT 2004)

24. If $x > 0$ and $\log_3 x + \log_3 \sqrt{x} + \log_3 (\sqrt[4]{x}) + \log_3 (\sqrt[8]{x})$

+ $\log_3 \sqrt[16]{x} + \dots = 4$, then x equals

(a) 1 (b) 9 (c) 27 (d) 81

(VITEEE 2007)

25. If a, b, c, d are in G.P, then $(a + b + c + d)^2$ is equal to

(a) $(a + b)^2 + (c + d)^2 + 2(b + c)^2$

(b) $(a + b)^2 + (c + d)^2 + 2(a + c)^2$

(c) $(a + b)^2 + (c + d)^2 + 2(b + d)^2$

(d) $(a + b)^2 + (c + d)^2 + (b + c)^2$

(Kerala PET 2012)

ANSWERS

1. (a) 2. (c) 3. (c) 4. (a) 5. (b) 6. (c) 7. (c) 8. (c) 9. (c) 10. (a)
11. (c) 12. (a) 13. (c) 14. (c) 15. (a) 16. (a) 17. (c) 18. (b) 19. (b) 20. (c)
21. (a) 22. (d) 23. (b) 24. (b) 25. (a)

HINTS AND SOLUTIONS

1. Here first term $a = 3$, common ratio $r = \frac{3\sqrt{3}}{3} = \sqrt{3}$

Let the n th term be 2187. Then,

$$T_n = ar^{n-1} = 2187$$

$$\Rightarrow 3 \times (\sqrt{3})^{n-1} = 2187$$

$$\Rightarrow (\sqrt{3})^{n-1} = 729$$

$$\Rightarrow (\sqrt{3})^{n-1} = 3^6 = (\sqrt{3})^{12}$$

$$\Rightarrow n-1 = 12 \Rightarrow n = 13.$$

2. Given, $n!$, $3 \times n!$ and $(n+1)!$ are in G.P

$$\Rightarrow (3 \times n!)^2 = n! \times (n+1)!$$

$$(\because a, b, c \text{ in G.P.} \Rightarrow b^2 = ac)$$

$$\Rightarrow 9 \times (n!)^2 = n! \times (n+1)!$$

$$\Rightarrow 9 \times n! = (n+1)!$$

$$\Rightarrow 9n! = (n+1) \cdot n! \quad (\because n! = 1.2.3. \dots n)$$

$$\Rightarrow n+1 = 9 \Rightarrow n = 8.$$

3. 1, $x, y, z, 16$ are in G.P.

$$\therefore a = 1, n = 5, T_5 = ar^4 = 16$$

$$\Rightarrow r^4 = 16 \Rightarrow r = 2$$

$$\therefore x = ar = 2, y = ar^2 = 4, z = ar^3 = 8$$

$$\therefore x + y + z = 2 + 4 + 8 = 14.$$

4. In the G.P, 1, 2, $2^2, \dots, 2^n$, there are $(n+1)$ terms.

$$\therefore \text{Geometric mean of this G.P.} = (1 \times 2 \times 2^2 \times \dots \times 2^n)^{\frac{1}{n+1}}$$

$$= (2^{1+2+3+\dots+n})^{\frac{1}{n+1}}$$

$$= \left(2^{\frac{n(n+1)}{2}}\right)^{\frac{1}{n+1}} = 2^{n/2}.$$

5. Let the first term and common ratio of the G.P. be a and r respectively.

$$\text{Then, } T_1 = a = 729, T_7 = ar^6 = 64$$

$$\therefore \frac{ar^6}{a} = \frac{64}{729} \Rightarrow r^6 = \frac{64}{729} = \frac{2^6}{3^6} \Rightarrow r = \frac{2}{3}.$$

$$\therefore S_7 = \frac{a(1-r^n)}{1-r} = \frac{729 \left(1 - \left(\frac{2}{3}\right)^7\right)}{1 - 2/3}$$

$$= \frac{729 \left(1 - \frac{128}{2187}\right)}{\frac{1}{3}} = 2187 \left(\frac{2187 - 128}{2187}\right) = 2059.$$

6. Let a and r be the first term and common ratio respectively of the given G.P. Then,

$$T_3 = ar^2 = 3$$

$$\text{Required product} = a \cdot ar \cdot ar^2 \cdot ar^3 \cdot ar^4$$

$$= a^5 r^{15} = (ar^3)^5 = 3^5 = 243.$$

7. Let a and r be the first term and common ratio respectively of the given G.P. Then, $t_2 = ar, t_4 = ar^3, t_5 = ar^4, t_6 = ar^5$

$$\text{Given, } t_2 + t_5 = ar + ar^4 = 216 \quad \dots(i)$$

$$\text{and } \frac{t_4}{t_6} = \frac{ar^3}{ar^5} = \frac{1}{4} \Rightarrow \frac{1}{r^2} = \frac{1}{4} \Rightarrow \frac{1}{r} = \frac{1}{2} \quad \dots(ii)$$

Now putting $r = 2$, in (i) we get

$$2a + 16a = 216 \Rightarrow 18a = 216 \Rightarrow a = 12.$$

8. Let $S_n = \frac{1}{2} + \frac{3}{4} + \frac{7}{8} + \frac{15}{16} + \dots$ to n terms

$$= \left(1 - \frac{1}{2}\right) + \left(1 - \frac{1}{4}\right) + \left(1 - \frac{1}{8}\right) + \left(1 - \frac{1}{16}\right) + \dots \text{ to } n \text{ terms}$$

$$= (1 + 1 + 1 + 1 + \dots \text{ to } n \text{ terms})$$

$$- \left\{ \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \frac{1}{16} + \dots \text{ to } n \text{ terms} \right\}$$

$$= n - \frac{1}{2} \left(1 - \left(\frac{1}{2} \right)^n \right) = n - (1 - 2^{-n}) = n + 2^{-n} = 1.$$

9. Let the G.P. be a, ar, ar^2, \dots

As all the terms of the given G.P. are positive, $a > 0, r > 0$.

Given, $a = ar + ar^2$

$$\Rightarrow ar^2 + ar - a = 0$$

$$\Rightarrow r^2 + r - 1 = 0.$$

$$\therefore r = \frac{-1 \pm \sqrt{1+4}}{2} = \frac{-1 \pm \sqrt{5}}{2}$$

$$\Rightarrow r = \frac{\sqrt{5}-1}{2}. \quad \left(\because \frac{-1-\sqrt{5}}{2} \text{ is a negative quantity and } r > 0 \right)$$

10. Let the given G.P. be $a, ar, ar^2, ar^3, \dots, \infty$.

As the given infinite G.P. has a finite sum, $|r| < 1$

Also, given $a = 2(ar + ar^2 + ar^3 + \dots + \infty)$

$$\Rightarrow a = 2 \left(\frac{ar}{1-r} \right) \left(\because S_{\infty} = \frac{a}{1-r}. \text{ Here } a = ar, r = r \right)$$

$$\Rightarrow a - ar = 2ar$$

$$\Rightarrow 1 - r = 2r \Rightarrow 3r = 1 \Rightarrow r = \frac{1}{3}.$$

11. Sum of an infinite G.P. = $\frac{a}{1-r}$, where first term = a ,

common ratio = r and $|r| < 1$

Given, $\frac{a}{1-r} = 4$ and $ar = \frac{3}{4}$

$$\Rightarrow a = 4 - 4r \text{ and } a = \frac{3}{4r}$$

$$\Rightarrow \frac{3}{4r} = 4 - 4r \Rightarrow 3 = 16r - 16r^2$$

$$\Rightarrow 16r^2 - 16r + 3 = 0$$

$$\Rightarrow (4r - 3)(4r - 1) = 0$$

$$\Rightarrow 4r = 3 \text{ or } 4r = 1 \Rightarrow r = \frac{3}{4} \text{ or } \frac{1}{4}$$

Now when $r = \frac{3}{4}, a = \frac{3}{4 \times \frac{3}{4}} = 1$

$$r = \frac{1}{4}, a = \frac{3}{4 \times \frac{1}{4}} = 3$$

$$\therefore (a, r) = \left(1, \frac{3}{4} \right) \text{ or } \left(3, \frac{1}{4} \right).$$

12. $x, 2x + 2, 3x + 3$, are in G.P.

$$\Rightarrow (2x + 2)^2 = x(3x + 3)$$

$$\Rightarrow 4x^2 + 8x + 4 = 3x^2 + 3x$$

$$\Rightarrow x^2 + 5x + 4 = 0$$

$$\Rightarrow (x + 4)(x + 1) = 0$$

$$\Rightarrow x = -1 \text{ or } -4$$

Let the fourth term of the given G.P. be a . Then,

$$r = \frac{a}{3x+3} = \frac{2x+2}{x}$$

$$\Rightarrow a = \frac{(2x+2)(3x+3)}{x}$$

When $x = -4$, Fourth term $a = \frac{(-8+2)(-24+3)}{-4}$

$$= \frac{-6 \times -21}{-4} = -\frac{27}{2}$$

When $x = -1$, $a = \frac{(-2+2)(-3+3)}{-1} = 0$.

$$\therefore \text{Fourth term} = -\frac{27}{2}.$$

13. Let the six numbers in G.P. be $\frac{a}{r^5}, \frac{a}{r^3}, \frac{a}{r}, ar, ar^3, ar^5$

Given, $\frac{a}{r^5} \times \frac{a}{r^3} \times \frac{a}{r} \times ar \times ar^3 \times ar^5 = 1000$

$$\Rightarrow a^6 = 1000 \Rightarrow a = \sqrt[3]{1000}$$

Given, $T_4 = ar = 1 \Rightarrow \sqrt[3]{1000}r = 1 \Rightarrow r = \frac{1}{\sqrt[3]{1000}}$

$$\therefore \text{Last term of G.P.} = ar^5 = \sqrt[3]{1000} \times \frac{1}{(\sqrt[3]{1000})^5} = \frac{1}{100}.$$

14. a is the A.M of b and $c \Rightarrow 2a = b + c$

Given, G_1 and G_2 are G.Ms between b and c

$\Rightarrow b, G_1, G_2, c$ are in G.P.

Let r be the common ratio of the G.P.

$$\Rightarrow G_1 = br, G_2 = br^2, c = br^3$$

Now $c = br^3 \Rightarrow r^3 = \frac{c}{b} \Rightarrow r = \left(\frac{c}{b} \right)^{\frac{1}{3}}$

$$\therefore G_1 = b \times \frac{c^{1/3}}{b^{1/3}} = b^{2/3} c^{1/3} = (b^2c)^{1/3}$$

$$G_2 = b \times \left(\frac{c}{b} \right)^{\frac{2}{3}} = \frac{b \times c^{2/3}}{b^{2/3}} = b^{1/3} c^{2/3} = (bc^2)^{1/3}$$

$$\begin{aligned} \text{Now } G_1^3 + G_2^3 &= \left((b^2c)^{1/3} \right)^3 + \left((bc^2)^{1/3} \right)^3 \\ &= b^2c + bc^2 = bc(b+c) = bc \cdot 2a = 2abc. \end{aligned}$$

15. a, b, c are in A.P. $\Rightarrow b - a = c - b \dots(i)$

$(b - a), (c - b), a$ are in G.P. $\Rightarrow (c - b)^2 = (b - a)a \dots(ii)$

\therefore From (i) and (ii)

$$\Rightarrow (b - a)^2 = (b - a)a$$

$$\Rightarrow (b - a)[(b - a) - a] = 0$$

$$\Rightarrow b - a = 0 \text{ or } b - 2a = 0$$

$$\Rightarrow b = 2a$$

$(\because a$ and b are distinct and $b - a \neq 0)$

a, b, c are in A.P. $\Rightarrow 2b = a + c$

$$\Rightarrow 4a = a + c \Rightarrow c = 3a.$$

$$\therefore a : b : c = a : 2a : 3a = 1 : 2 : 3.$$

$$\begin{aligned}
 16. \text{ Let } S_{100} &= 9 + 99 + 999 + \dots \text{ upto 100 terms} \\
 &= (10 - 1) + (100 - 1) + (1000 - 1) \\
 &\quad + \dots + \text{ upto 100 terms} \\
 &= (10 + 10^2 + 10^3 + \dots \text{ upto 100 terms}) \\
 &\quad - (1 + 1 + 1 + \dots \text{ upto 100 terms}) \\
 &= \frac{10(10^{100} - 1)}{10 - 1} - 100 \\
 &\quad \left(\because S_n = \frac{a(r^n - 1)}{r - 1} \text{ when } r > 1 \right) \\
 &= \frac{10}{9}(10^{100} - 1) - 100.
 \end{aligned}$$

$$\begin{aligned}
 17. \text{ Let } S_{20} &= 0.7 + 0.77 + 0.777 + \dots \text{ upto 20 terms} \\
 &= 7(0.1 + 0.11 + 0.111 + \dots \text{ upto 20 terms}) \\
 &= \frac{7}{9}(0.9 + 0.99 + 0.999 + \dots \text{ upto 20 terms}) \\
 &= \frac{7}{9}[(1 - 0.1) + (1 - 0.01) + (1 - 0.001) + \dots \\
 &\quad \text{20 terms}] \\
 &= \frac{7}{9}[(1 + 1 + 1 + \dots \text{ upto 20 terms}) - (0.1 \\
 &\quad + 0.01 + 0.001 + \dots \text{ upto 20 terms})] \\
 &= \frac{7}{9}\left[20 - \frac{0.1\{1 - (0.1)^{20}\}}{(1 - 0.1)}\right] \\
 &\quad \left(\because S_n = \frac{a(1 - r^n)}{1 - r}, \text{ when } r < 1 \right) \\
 &= \frac{7}{9}\left[20 - \frac{1}{9}\left(1 - \left(\frac{1}{10}\right)^{20}\right)\right] = \frac{7}{9}\left[20 - \frac{1}{9} + \frac{10^{-20}}{9}\right] \\
 &= \frac{7}{9}\left[\frac{179 + 10^{-20}}{9}\right] = \frac{7}{81}(179 + 10^{-20}).
 \end{aligned}$$

$$\begin{aligned}
 18. \text{ Given } &\left(x - \frac{1}{y}\right) + \left(x^2 - \frac{1}{y^2}\right) + \dots \text{ to } \infty = \frac{2}{3} \\
 \Rightarrow &(x + x^2 + x^3 + \dots \text{ to } \infty) - \left(\frac{1}{y} + \frac{1}{y^2} + \dots \text{ to } \infty\right) = \frac{2}{3} \\
 \Rightarrow &\frac{x}{1 - x} - \frac{\frac{1}{y}}{1 - \frac{1}{y}} = \frac{2}{3} \Rightarrow \frac{x}{1 - x} - \frac{1}{y - 1} = \frac{2}{3} \\
 \Rightarrow &\frac{\frac{2}{y}}{1 - \frac{2}{y}} - \frac{1}{y - 1} = \frac{2}{3} \quad (\because xy = 2 \Rightarrow x = 2/y) \\
 \Rightarrow &\frac{2}{y - 2} - \frac{1}{y - 1} = \frac{2}{3} \Rightarrow \frac{2y - 2 - y + 2}{y^2 - 3y + 2} = \frac{2}{3} \\
 \Rightarrow &3y = 2(y^2 - 3y + 2) \\
 \Rightarrow &2y^2 - 9y + 4 = 0 \Rightarrow 2y^2 - 8y - y + 4 = 0 \\
 \Rightarrow &(2y - 1)(y - 4) = 0.
 \end{aligned}$$

$$\Rightarrow y = \frac{1}{2} \text{ or } y = 4$$

$$\text{When } y = \frac{1}{2}, x = 4 \text{ and } y = 4, x = \frac{1}{2}$$

$$\therefore x < 1 \quad \therefore y = 4, x = \frac{1}{2} \text{ is true.}$$

19. Let a and r be the first term and common ratio respectively of the given G.P.

$$\text{Then } \begin{cases} a + ar = 12 & \dots(i) \\ ar^2 + ar^3 = 48 & \dots(ii) \end{cases}$$

$$\Rightarrow \frac{ar^2(1+r)}{a(1+r)} = \frac{48}{12} \quad (\text{Dividing (ii) by (i)})$$

$$\Rightarrow r^2 = 4 \Rightarrow r = 2 \Rightarrow r = -2$$

as the terms of the G.P. are alternately positive and negative.

$$\text{Now } a(1+r) = 12 \Rightarrow a(1-2) = 12 \Rightarrow a = -12.$$

20. Let a and r be the first term and common ratio respectively of the given G.P.

$$P\text{th term of G.P.} = ar^{P-1} = 64 = 2^6 \quad \dots(i)$$

$$Q\text{th term of G.P.} = ar^{Q-1} = 27 = 3^3 \quad \dots(ii)$$

$$R\text{th term of G.P.} = ar^{R-1} = 36 = 2^2 \cdot 3^2 \quad \dots(iii)$$

$$\text{Now from (i), } 2 = a^{1/6} r^{(P-1)/6} \quad \dots(iv)$$

$$\text{From (ii), } 3 = a^{1/3} r^{(Q-1)/3} \quad \dots(v)$$

$$\text{From (iii), } 2.3 = a^{1/2} r^{(R-1)/2} \quad \dots(vi)$$

\therefore From (iv), (v) and (vi) we have

$$\Rightarrow a^{1/6} r^{P/6} \cdot a^{1/3} r^{Q/3} = a^{1/2} r^{R/2}$$

$$\Rightarrow a^{1/6 + 1/3} r^{P/6 + Q/3} = a^{1/2} r^{R/2}$$

$$\Rightarrow a^{1/2} r^{P-1+2Q-2} = a^{1/2} r^{R-1}$$

$$\Rightarrow \frac{P-1+2Q-2}{6} = \frac{R-1}{2}$$

$$\Rightarrow P + 2Q - 3 = 3R - 3$$

$$\Rightarrow P + 2Q = 3R.$$

21. Let the first term of the A.P. be a and common difference d .

$$\text{Given } d = 2 \Rightarrow T_{11} \text{ of A.P.} = a + 10d = a + 20.$$

Let the first term of the G.P. be b and common ratio r .

$$\text{Given } r = 2.$$

Now, the middle term of A.P. = middle term of G.P.

$$\Rightarrow T_6 \text{ of A.P.} = a + 5d = T_6 \text{ of G.P.} = br^5$$

$$\Rightarrow a + 5d = br^5$$

$$\Rightarrow a + 10 = 32b \quad (\because r = 2) \quad \dots(i)$$

Also the last term of A.P. is the first term of G.P.

$$\therefore b = T_{11} \text{ of A.P.} = a + 20 \quad \dots(ii)$$

\therefore From (i) and (ii)

$$a + 10 = 32.(a + 20)$$

$$\Rightarrow 31a = -630 \Rightarrow a = \frac{-630}{31}$$

$$\begin{aligned} \therefore \text{Middle term of the entire sequence of 21 terms} &= 11\text{th term} \\ &= a + 10d \\ &= \frac{-630}{31} + 20 = \frac{-630 + 620}{31} = \frac{-10}{31}. \end{aligned}$$

22. Since a and b are proper fractions, $|a| < 1, |b| < 1$

$$\therefore x = 1 + a + a^2 + \dots \infty = \frac{1}{1-a} \quad \left(\because S_\infty = \frac{a}{1-r} \right)$$

and $y = 1 + b + b^2 + \dots \infty = \frac{1}{1-b}$

Also, $1 + ab + a^2b^2 + \dots \infty = \frac{1}{1-ab} \quad \dots(i)$

Now $x = \frac{1}{1-a} \Rightarrow x - xa = 1$

$$\Rightarrow xa = x - 1 \Rightarrow a = \frac{x-1}{x} \quad \dots(ii)$$

$$y = \frac{1}{1-b} \Rightarrow y - yb = 1$$

$$\Rightarrow yb = y - 1 \Rightarrow b = \frac{y-1}{y} \quad \dots(iii)$$

\therefore Putting the values of a & b from (ii) and (iii) in (i), we get

$$\begin{aligned} \text{Reqd. sum} &= \frac{1}{1-ab} = \frac{1}{1 - \left(\frac{x-1}{x}\right)\left(\frac{y-1}{y}\right)} \\ &= \frac{xy}{xy - (xy - x - y + 1)} = \frac{xy}{x + y - 1}. \end{aligned}$$

23. Let the common ratio of the given G.P be r . Then,

$$S_\infty = \frac{x}{1-r} \Rightarrow 5 = \frac{x}{1-r} \Rightarrow 5 - 5r = x$$

$$\Rightarrow r = \frac{5-x}{5} = 1 - \frac{x}{5}$$

\therefore Sum to infinity of the given series is a finite quantity, $|r| < 1$.

$$\therefore \left| 1 - \frac{x}{5} \right| < 1 \Rightarrow -1 < \left(1 - \frac{x}{5} \right) < 1$$

$$\Rightarrow -1 < (x/5 - 1) < 1$$

(Multiplying the inequality by (-1))

$$\Rightarrow 0 < \frac{x}{5} < 2 \Rightarrow 0 < x < 10.$$

24. $\log_3 x + \log_3 \sqrt{x} + \log_3 \sqrt[4]{x} + \log_3 \sqrt[8]{x} + \dots \infty = 4$

$$\Rightarrow \log_3 x + \log_3 x^{1/2} + \log_3 x^{1/4} + \log_3 (x^{1/8}) + \dots \infty = 4$$

$$\Rightarrow \log_3 (x \cdot x^{1/2} \cdot x^{1/4} \cdot x^{1/8} \dots \infty) = 4$$

$$\Rightarrow \log_3 (x^{1 + 1/2 + 1/4 + 1/8} \dots \infty) = 4$$

$$\Rightarrow \log_3 x^{\frac{1}{1-1/2}} = 4$$

($\because S_\infty = a/1-r$)

$$\Rightarrow \log_3 x^2 = 4 \Rightarrow x^2 = 3^4 = (3^2)^2 \Rightarrow x = 9.$$

25. a, b, c, d are in G.P.

$\Rightarrow b = ar, c = ar^2, d = ar^3$, where r is the common ratio of the G.P.

$$\begin{aligned} \text{LHS} &= (a + b + c + d)^2 = (a + ar + ar^2 + ar^3)^2 \\ &= \{a(1 + r + r^2 + r^3)\}^2 = \{a((1+r) + r^2(1+r))\}^2 \\ &= \{a(1+r)(1+r^2)\}^2 = a^2(1+r)^2(1+r^2)^2 \\ &= a^2(1+r)^2(r^4 + 2r^2 + 1) \\ &= a^2(r^4(1+r)^2 + 2r^2(1+r)^2 + (1+r)^2) \\ &= a^2r^4(1+r)^2 + 2a^2r^2(1+r)^2 + a^2(1+r)^2 \\ &= (ar^2 + ar^3)^2 + 2(ar + ar^2)^2 + (a + ar)^2 \\ &= (c + d)^2 + 2(b + c)^2 + (a + b)^2. \end{aligned}$$

HARMONIC PROGRESSION (H.P.)

KEY FACTS

1. A sequence of numbers is said to form a **Harmonic Progression**, when the reciprocals of the numbers form an **Arithmetic progression**.

Ex. (i) $1 + 2 + 3 + 4 + \dots$ is an A.P

$$\Rightarrow 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \dots \text{ is an H.P.}$$

(ii) $a + (a + d) + (a + 2d) + (a + 3d) \dots$ is an A.P.

$$\Rightarrow \frac{1}{a} + \frac{1}{(a+d)} + \frac{1}{(a+2d)} + \frac{1}{(a+3d)} + \dots \text{ is an H.P.}$$

2. **n th term of an H.P.**

The n th term of an H.P. is the reciprocal of the n th term of the A.P. formed by the reciprocals of the terms of the H.P.

If the given H.P. is $\frac{1}{a}, \frac{1}{a+d}, \frac{1}{a+2d}, \dots$, then its n th term is $\frac{1}{a + (n-1)d}$.

Ex. Find the 7th term of the H.P. $\frac{2}{13}, \frac{1}{6}, \frac{2}{11}, \dots$.

The reciprocals of the terms of the given H.P, i.e., $\frac{13}{2}, 6, \frac{11}{2}, \dots$ form an A.P with first term = $\frac{13}{2}$ and common difference = $6 - \frac{13}{2} = -\frac{1}{2}$.

$$\therefore \text{7th term of this A.P.} = a + 6d = \frac{13}{2} + \left(-\frac{1}{2}\right) \times 6 = \frac{7}{2}.$$

$$\text{Hence 7th of the given H.P.} = \frac{1}{7/2} = \frac{2}{7}.$$

3. Harmonic Mean

If a, H, b are three quantities in H.P., then H is said to be the **Harmonic Mean** between a and b .
 a, H, b are in H.P

$$\Rightarrow \frac{1}{a}, \frac{1}{H}, \frac{1}{b} \text{ are in A.P.}$$

$$\Rightarrow \frac{1}{H} - \frac{1}{a} = \frac{1}{b} - \frac{1}{H} \Rightarrow \frac{2}{H} = \frac{1}{b} + \frac{1}{a} \Rightarrow \frac{2}{H} = \frac{a+b}{ab} \Rightarrow H = \frac{2ab}{a+b}.$$

Ex. The **Harmonic Mean** between 3 and -5 is:

$$H = \frac{2 \times 3 \times (-5)}{3 + (-5)} = \frac{-30}{-2} = 15.$$

4. If $a_1, a_2, a_3, a_4, \dots, a_n$ are n non-zero numbers in H.P., then their

$$\text{Harmonic Mean} = \frac{\left(\frac{1}{a_1} + \frac{1}{a_2} + \frac{1}{a_3} + \dots + \frac{1}{a_n}\right)}{n}$$

5. Relations between the three series, i.e., A.P, G.P and H.P.

(a) Three numbers a, b, c will be in A.P, G.P or in H.P. according as $\frac{a-b}{b-c} = \frac{a}{a}, \frac{a-b}{b-c} = \frac{a}{b}, \frac{a-b}{b-c} = \frac{a}{c}$ respectively.

$$(i) \frac{a-b}{b-c} = \frac{a}{a} \Rightarrow a^2 - ba = ab - ac \Rightarrow 2ab = a^2 + ac \Rightarrow 2b = a + c \Rightarrow a, b, c \text{ are in A.P.}$$

$$(ii) \frac{a-b}{b-c} = \frac{a}{b} \Rightarrow ab - b^2 = ab - ca \Rightarrow b^2 = ac \Rightarrow a, b, c \text{ are in G.P.}$$

$$(iii) \frac{a-b}{b-c} = \frac{a}{c} \Rightarrow ac - bc = ab - ac \Rightarrow 2ac = ab + bc \Rightarrow b = \frac{2ac}{a+c} \Rightarrow a, b, c \text{ are H.P.}$$

(b) If A, G, H are respectively the Arithmetic, the Geometric and the Harmonic means between any two unequal positive numbers, then

(i) A, G, H are in G.P (ii) $A > G > H$.

(i) Let the two positive numbers be a and b . Then,

$$A = \frac{a+b}{2}, G = \sqrt{ab}, H = \frac{2ab}{a+b}$$

$$\text{Then } A \times H = \left(\frac{a+b}{2}\right) \times \frac{2ab}{a+b} = ab = G^2$$

$$\Rightarrow G = \sqrt{AH} \Rightarrow A, G, H \text{ are in G.P.}$$

$$(ii) \quad A - G = \frac{a+b}{2} - \sqrt{ab} = \frac{a+b-2\sqrt{ab}}{2} = \frac{(\sqrt{a}-\sqrt{b})^2}{2} > 0 \Rightarrow A - G > 0 \Rightarrow A > G$$

$$\text{Also } G - H = \sqrt{ab} - \frac{2ab}{a+b} = \frac{(a+b)\sqrt{ab} - 2ab}{a+b} = \frac{\sqrt{ab}(a+b-2\sqrt{ab})}{a+b} = \frac{\sqrt{ab}}{a+b}(\sqrt{a}-\sqrt{b})^2 > 0$$

$$\Rightarrow G - H > 0 \Rightarrow G > H$$

$$\therefore A > G > H.$$

SOLVED EXAMPLES

Ex. 1. The third and seventh terms of an H.P are $\frac{7}{11}$ and $\frac{7}{31}$. Find the first term and the eighth term.

Sol. Third and seventh terms of an H.P are $\frac{7}{11}$ and $\frac{7}{31}$

\Rightarrow Third and seventh terms of an A.P are $\frac{11}{7}$ and $\frac{31}{7}$

Let the first term of the A.P. be a and common difference d .

$$\text{Then, } T_3 = a + 2d = \frac{11}{7} \quad \dots(i)$$

$$T_7 = a + 6d = \frac{31}{7} \quad \dots(ii)$$

$$(ii) - (i) \Rightarrow 4d = \frac{20}{7} \Rightarrow d = \frac{5}{7}$$

$$\therefore \text{ From (i) } a + \frac{10}{7} = \frac{11}{7} \Rightarrow a = \frac{1}{7}$$

$$T_8 = a + 7d = \frac{1}{7} + 7 \times \frac{5}{7} = \frac{1}{7} + 5 = \frac{36}{7}$$

\therefore The first and eighth terms of the H.P. are respectively the reciprocals of the first and eighth terms of the A.P, i.e. 7 and $\frac{7}{36}$.

Ex. 2. Insert three harmonic means between 5 and 6.

Sol. 3 harmonic means between 5 and 6

\Rightarrow 3 arithmetic means between $\frac{1}{5}$ and $\frac{1}{6}$

Let A_1, A_2, A_3 be the arithmetic means between $\frac{1}{5}$ and $\frac{1}{6}$.

Then, $\frac{1}{5}, A_1, A_2, A_3, \frac{1}{6}$ form an A.P,

where $t_1 = a = \frac{1}{5}$, $t_5 = a + 4d = \frac{1}{6}$

$$\therefore \frac{1}{5} + 4d = \frac{1}{6} \Rightarrow 4d = \frac{1}{6} - \frac{1}{5} = -\frac{1}{30} \Rightarrow d = -\frac{1}{120}$$

$$\therefore A_1 = a + d = \frac{1}{5} + \left(-\frac{1}{120}\right) = \frac{23}{120}$$

$$A_2 = a + 2d = \frac{1}{5} + 2 \times \left(-\frac{1}{120}\right) = \frac{1}{5} - \frac{1}{60} = \frac{11}{60}$$

$$A_3 = a + 3d = \frac{1}{5} + 3 \times \left(-\frac{1}{120} \right) = \frac{1}{5} - \frac{1}{40} = \frac{7}{40}$$

∴ Required harmonic means are $\frac{120}{23}, \frac{60}{11}, \frac{40}{7}$.

Ex. 3. If the m th term of an H.P. is n and n th term is m , show that the r th term is $\frac{mn}{r}$.

Sol. Let the corresponding A.P. be $a, a + d, a + 2d, \dots$

Since the m th term and n th term of the H.P. are n and m respectively, then for the A.P.,

$$m\text{th term} = a + (m - 1)d = \frac{1}{n} \quad \dots(i)$$

$$n\text{th term} = a + (n - 1)d = \frac{1}{m} \quad \dots(ii)$$

$$(ii) - (i) \Rightarrow (n - 1)d - (m - 1)d = \frac{1}{m} - \frac{1}{n}$$

$$\Rightarrow (n - m)d = \frac{n - m}{mn} \Rightarrow d = \frac{1}{mn}$$

$$\text{Putting in (i), } a + (m - 1) \times \frac{1}{mn} = \frac{1}{n}$$

$$\Rightarrow a = \frac{1}{n} - \frac{1}{n} + \frac{1}{mn} = \frac{1}{mn}$$

$$\therefore t_r = a + (r - 1)d = \frac{1}{mn} + (r - 1) \frac{1}{mn} = \frac{r}{mn}$$

∴ r th term of H.P. = $\frac{mn}{r}$.

Ex. 4. If H be the harmonic mean between x and y , then prove that $\frac{H + x}{H - x} + \frac{H + y}{H - y} = 2$.

Sol. H being the H.M. between x and y

$$\Rightarrow H = \frac{2xy}{x + y} \Rightarrow \frac{H}{x} = \frac{2y}{x + y} \text{ and } \frac{H}{y} = \frac{2x}{x + y}$$

$$\Rightarrow \frac{H + x}{H - x} = \frac{2y + x + y}{2y - (x + y)} \text{ and } \frac{H + y}{H - y} = \frac{2x + x + y}{2x - (x + y)}$$

(Using Componendo and Dividendo)

$$\Rightarrow \frac{H + x}{H - x} = \frac{3y + x}{y - x} \text{ and } \frac{H + y}{H - y} = \frac{3x + y}{x - y}$$

$$\therefore \frac{H + x}{H - x} + \frac{H + y}{H - y} = \frac{3y + x}{y - x} + \frac{3x + y}{x - y} = \frac{3y + x - 3x - y}{y - x} = \frac{2(y - x)}{y - x} = 2.$$

Ex. 5. In an H.P., p th term is qr and q th term is pr , show that r th term is pq .

$$\text{Sol. } T_p \text{ of H.P.} = qr \Rightarrow T_p \text{ of A.P.} = \frac{1}{qr} \Rightarrow a + (p - 1)d = \frac{1}{qr} \quad \dots(i)$$

Where a and d are the first term and common difference respectively of the A.P.

$$\text{Also } T_q \text{ of H.P.} = pr \Rightarrow T_q \text{ of A.P.} = \frac{1}{pr} \Rightarrow a + (q - 1)d = \frac{1}{pr} \quad \dots(ii)$$

$$\begin{aligned} \text{Eqn (ii)} - \text{Eqn (i)} &\Rightarrow (q-1)d - (p-1)d = \frac{1}{pr} - \frac{1}{qr} \Rightarrow (q-p)d = \frac{q-p}{pqr} \Rightarrow d = \frac{1}{pqr} \\ \Rightarrow a + (p-1)\frac{1}{pqr} &= \frac{1}{qr} \Rightarrow a = \frac{1}{qr} - (p-1)\frac{1}{pqr} = \frac{1}{qr} - \frac{1}{qr} + \frac{1}{pqr} = \frac{1}{pqr} \\ \therefore T_r \text{ of A.P} &= a + (r-1)d = \frac{1}{pqr} + (r-1)\frac{1}{pqr} = \frac{1}{pqr} + \frac{1}{pq} - \frac{1}{pqr} = \frac{1}{pq} \\ \Rightarrow T_r \text{ of H.P} &= pq. \end{aligned}$$

Ex. 6. Let a, b, c be in A.P and $|a| < 1, |b| < 1$ and $|c| < 1$. If

$$x = 1 + a + a^2 + \dots \text{ to } \infty$$

$$y = 1 + b + b^2 + \dots \text{ to } \infty$$

$$z = 1 + c + c^2 + \dots \text{ to } \infty, \text{ then show that } x, y, z \text{ are in H.P.}$$

Sol.

$$\begin{aligned} x = 1 + a + a^2 + \dots \text{ to } \infty &= \frac{1}{1-a} \\ y = 1 + b + b^2 + \dots \text{ to } \infty &= \frac{1}{1-b} \\ z = 1 + c + c^2 + \dots \text{ to } \infty &= \frac{1}{1-c}. \end{aligned}$$

Now, a, b, c are in A.P.

$$\Rightarrow 1-a, 1-b, 1-c \text{ are in A.P.}$$

$$\Rightarrow \frac{1}{1-a}, \frac{1}{1-b}, \frac{1}{1-c} \text{ are in H.P.}$$

$$\Rightarrow x, y, z \text{ are in H.P.}$$

Ex. 7. If $a_1, a_2, a_3, \dots, a_n$ are in H.P. then what will $(a_1a_2 + a_2a_3 + \dots + a_{n-1}a_n)$ equal to?

(AMU 2003)

Sol. $a_1, a_2, a_3, \dots, a_n$ are in H.P.

$$\Rightarrow \frac{1}{a_1}, \frac{1}{a_2}, \frac{1}{a_3}, \dots, \frac{1}{a_n} \text{ are in A.P}$$

If d is the common difference of the A.P., then,

$$\frac{1}{a_2} - \frac{1}{a_1} = d, \frac{1}{a_3} - \frac{1}{a_2} = d, \dots, \frac{1}{a_n} - \frac{1}{a_{n-1}} = d \Rightarrow \frac{a_1 - a_2}{a_1a_2} = d, \frac{a_2 - a_3}{a_2a_3} = d, \dots, \frac{a_{n-1} - a_n}{a_{n-1}a_n} = d$$

$$\Rightarrow a_1 - a_2 = a_1a_2d, a_2 - a_3 = a_2a_3d, \dots, a_{n-1} - a_n = a_{n-1}a_nd$$

$$\Rightarrow (a_1 - a_2 + a_2 - a_3 + \dots + a_{n-1} - a_n) = a_1a_2d + a_2a_3d + \dots + a_{n-1}a_nd$$

$$\Rightarrow (a_1 - a_n) = (a_1a_2 + a_2a_3 + \dots + a_{n-1}a_n)d \tag{... (i)}$$

Also, $\frac{1}{a_1}, \frac{1}{a_2}, \dots, \frac{1}{a_n}$ is an A.P with common difference d

$$\Rightarrow \frac{1}{a_n} = \frac{1}{a_1} + (n-1)d \Rightarrow (n-1)d = \frac{1}{a_n} - \frac{1}{a_1} \Rightarrow (n-1)d = \frac{a_1 - a_n}{a_1a_n}$$

$$\Rightarrow a_1 - a_n = (n-1)d a_1a_n \tag{... (ii)}$$

\therefore From (i) and (ii)

$$(n-1)d a_1a_n = (a_1a_2 + a_2a_3 + \dots + a_{n-1}a_n) d$$

$$\Rightarrow (a_1a_2 + a_2a_3 + \dots + a_{n-1}a_n) = (n-1) a_1a_n.$$

Ex. 8. If $\frac{1}{a^2}, \frac{1}{b^2}, \frac{1}{c^2}$ are in H.P, show that $b + c, c + a, a + b$ are in H.P.

Sol. We know by the property of H.P, if x, y, z are H.P, $\frac{x-y}{y-z} = \frac{x}{z}$

$$\begin{aligned} \therefore \frac{1}{a^2}, \frac{1}{b^2}, \frac{1}{c^2} \text{ are in H.P.} &\Rightarrow \frac{\frac{1}{a^2} - \frac{1}{b^2}}{\frac{1}{b^2} - \frac{1}{c^2}} = \frac{\frac{1}{a^2}}{\frac{1}{c^2}} \Rightarrow \frac{\frac{b^2 - a^2}{a^2 b^2}}{\frac{c^2 - b^2}{b^2 c^2}} = \frac{c^2}{a^2} \Rightarrow \frac{c^2 (b^2 - a^2)}{a^2 (c^2 - b^2)} = \frac{c^2}{a^2} \\ \Rightarrow & b^2 - a^2 = c^2 - b^2 \Rightarrow 2b^2 = c^2 + a^2 \quad \dots(i) \end{aligned}$$

Now, for $b + c, c + a, a + b$ to be in H.P

$$\begin{aligned} \frac{(b+c) - (c+a)}{(c+a) - (a+b)} = \frac{b+c}{a+b} &\Rightarrow \frac{b-a}{c-b} = \frac{b+c}{a+b} \\ \Rightarrow (b-a)(b+a) &= (c+b)(c-b) \\ \Rightarrow b^2 - a^2 &= c^2 - b^2 \quad \text{or} \quad 2b^2 = c^2 + a^2, \text{ which is true from (i)} \\ \therefore b+c, c+a, a+b &\text{ are in H.P.} \end{aligned}$$

Ex. 9. The sum of the reciprocals of three numbers in H.P. is 15 and the product of the numbers is $\frac{1}{105}$. Find the numbers.

Sol. Let the three numbers in H.P. be $\frac{1}{a-d}, \frac{1}{a}, \frac{1}{a+d}$

$$\therefore (a-d) + a + (a+d) = 15$$

$$\Rightarrow 3a = 15 \Rightarrow a = 5$$

$$\text{Also } \frac{1}{(a-d)} \cdot \frac{1}{a} \cdot \frac{1}{(a+d)} = \frac{1}{105} \Rightarrow \frac{1}{a(a^2 - d^2)} = \frac{1}{105} \Rightarrow \frac{1}{5(25 - d^2)} = \frac{1}{105}$$

$$\Rightarrow 25 - d^2 = 21$$

$$\Rightarrow d^2 = 4 \Rightarrow d = \pm 2$$

Hence the numbers are $\frac{1}{3}, \frac{1}{5}, \frac{1}{7}$ when $d = 2$

or $\frac{1}{7}, \frac{1}{5}, \frac{1}{3}$ when $d = -2$.

Ex. 10. Insert between $\frac{1}{6}$ and $\frac{1}{16}$ two numbers such that the first three may be in H.P. and the last three in G.P.

Sol. Let a and b be the two numbers between $\frac{1}{6}$ and $\frac{1}{16}$. Then $\frac{1}{6}, a, b$ are in H.P.

$$\Rightarrow a = \frac{2 \times b \times \frac{1}{6}}{\frac{1}{6} + b} \quad \left(\because a, b, c \text{ in H.P.} \Rightarrow b = \frac{2ac}{a+c} \right)$$

$$\Rightarrow \frac{a}{6} + ab = \frac{b}{3} \Rightarrow a + 6ab = 2b \quad \dots(i)$$

Also, $a, b, \frac{1}{16}$ are in G.P.

$$\Rightarrow b^2 = a \times \frac{1}{16} \Rightarrow b^2 = \frac{a}{16} \Rightarrow a = 16b^2 \quad \dots(ii)$$

\therefore From (i) and (ii)

$$\Rightarrow 16b^2 + 6 \times 16b^2 \times b = 2b \Rightarrow 16b^2 + 96b^3 - 2b = 0 \Rightarrow b(16b + 96b^2 - 2) = 0$$

$$\Rightarrow b(96b^2 + 16b - 2) = 0 \Rightarrow b(48b^2 + 8b - 1) = 0$$

$$\Rightarrow b(12b - 1)(4b + 1) = 0$$

$$\Rightarrow b = \frac{1}{12} \text{ or } -\frac{1}{4} \text{ or } 0$$

Since $-\frac{1}{4}$ and 0 do not lie between $\frac{1}{6}$ and $\frac{1}{16}$, therefore, $b = \frac{1}{12}$

$$\text{Hence } a = 16b^2 = 16 \times \frac{1}{144} = \frac{1}{9}$$

\therefore The required numbers are $\frac{1}{4}$ and $\frac{1}{12}$.

Ex. 11. If A.M. between two numbers is to their G.M. as 5 : 4 and the difference of their G.M. and H.M. is 16/5, find the numbers.

Sol. Let the two numbers be a and b . Then,

$$\text{A.M.} = \frac{a+b}{2}$$

$$\text{G.M.} = \sqrt{ab}$$

$$\text{H.M.} = \frac{2ab}{a+b}$$

Given, A.M. : G.M. = 5 : 4

$$\Rightarrow \frac{(a+b)/2}{\sqrt{ab}} = \frac{5}{4} \Rightarrow \frac{a+b}{2\sqrt{ab}} = \frac{5}{4} \quad \dots(i)$$

Also, G.M. - H.M. = 16/5

$$\Rightarrow \sqrt{ab} - \frac{2ab}{a+b} = \frac{16}{5} \quad \dots(ii)$$

$$\text{From (i), } a+b = \frac{5 \times 2}{4} \sqrt{ab} \Rightarrow a+b = \frac{5}{2} \sqrt{ab}. \quad \dots(iii)$$

Putting this value of $(a+b)$ in (ii), we have

$$\sqrt{ab} - \frac{2ab}{\sqrt{ab}} \times \frac{2}{5} = \frac{16}{5} \Rightarrow \sqrt{ab} - \frac{4}{5} \sqrt{ab} = \frac{16}{5}$$

$$\Rightarrow \frac{1}{5} \sqrt{ab} = \frac{16}{5} \Rightarrow \sqrt{ab} = 16 \Rightarrow ab = 256$$

$$\therefore \text{ From (iii), } a+b = \frac{5}{2} \times 16 = 40.$$

$$\therefore (a-b)^2 = (a+b)^2 - 4ab = 40^2 - 4 \times 256 = 1600 - 1024 = 576$$

$$\Rightarrow a-b = \pm 24$$

Now solving $a+b = 40$ and $a-b = \pm 24$, we get

$$a = 32, b = 8 \quad \text{or} \quad a = 8, b = 32.$$

\therefore The numbers are **8 and 32**.

Ex. 12. If $\frac{a-x}{px} = \frac{a-y}{qy} = \frac{a-z}{rz}$ and p, q, r are in A.P., show that x, y, z are in H.P.

Sol. Let
$$\frac{a-x}{px} = \frac{a-y}{qy} = \frac{a-z}{rz} = k$$

Then,
$$\frac{a-x}{px} = k \Rightarrow \frac{a-x}{kx} = p \Rightarrow p = \frac{1}{k} \left(\frac{a}{x} - 1 \right)$$

Similarly,
$$q = \frac{1}{k} \left(\frac{a}{y} - 1 \right), r = \frac{1}{k} \left(\frac{a}{z} - 1 \right)$$

Now, p, q, r are in A.P.

$$\Rightarrow \frac{1}{k} \left(\frac{a}{x} - 1 \right), \frac{1}{k} \left(\frac{a}{y} - 1 \right), \frac{1}{k} \left(\frac{a}{z} - 1 \right) \text{ are in A.P.}$$

$$\Rightarrow \frac{a}{x} - 1, \frac{a}{y} - 1, \frac{a}{z} - 1 \text{ are in A.P.}$$

(Multiplying each term by k)

$$\Rightarrow \frac{a}{x}, \frac{a}{y}, \frac{a}{z} \text{ are in A.P.}$$

(Adding 1 to each term)

$$\Rightarrow \frac{1}{x}, \frac{1}{y}, \frac{1}{z} \text{ are in A.P.}$$

(Dividing each term by a)

$$\Rightarrow x, y, z \text{ are in H.P.}$$

Ex. 13. Let a, b be two positive real numbers. If a, A_1, A_2, b are in arithmetic progression, a, G_1, G_2, b are in geometric progression and a, H_1, H_2, b are in harmonic progression, then show that

$$\frac{G_1 G_2}{H_1 H_2} = \frac{A_1 + A_2}{H_1 + H_2} = \frac{(2a+b)(a+2b)}{9ab}$$

Sol. a, A_1, A_2, b are in A.P. $\Rightarrow A_1 - a = b - A_2 \Rightarrow A_1 + A_2 = a + b$... (i)

a, G_1, G_2, b are in G.P. $\Rightarrow \frac{G_1}{a} = \frac{b}{G_2} \Rightarrow G_1 G_2 = ab$... (ii)

a, H_1, H_2, b are in H.P.

$$\Rightarrow \frac{1}{a}, \frac{1}{H_1}, \frac{1}{H_2}, \frac{1}{b} \text{ are in A.P.}$$

$$\Rightarrow \frac{1}{H_1} - \frac{1}{a} = \frac{1}{b} - \frac{1}{H_2}$$

$$\Rightarrow \frac{1}{H_1} + \frac{1}{H_2} = \frac{1}{a} + \frac{1}{b}$$

$$\Rightarrow \frac{H_1 + H_2}{H_1 H_2} = \frac{a+b}{ab} = \frac{A_1 + A_2}{G_1 G_2}$$

(From (i) and (ii))

$$\Rightarrow \frac{G_1 G_2}{H_1 H_2} = \frac{A_1 + A_2}{H_1 + H_2}. \text{ Hence proved.}$$

Now $\frac{1}{a}, \frac{1}{H_1}, \frac{1}{H_2}$ are in A.P.

($\because a, H_1, H_2$, are in H.P.)

$$\Rightarrow \frac{1}{H_1} - \frac{1}{a} = \frac{1}{H_2} - \frac{1}{H_1}$$

$$\Rightarrow \frac{2}{H_1} - \frac{1}{H_2} = \frac{1}{a} \quad \dots(iii)$$

Also, $\frac{1}{H_1}, \frac{1}{H_2}, \frac{1}{b}$ are in A.P. ($\because H_1, H_2, b$ are in H.P.)

$$\Rightarrow \frac{1}{H_2} - \frac{1}{H_1} = \frac{1}{b} - \frac{1}{H_2} \Rightarrow \frac{2}{H_2} - \frac{1}{H_1} = \frac{1}{b} \quad \dots(iv)$$

$$\text{Eq. (iii)} + 2 \times \text{Eqn. (iv)} \Rightarrow \left(\frac{2}{H_1} - \frac{1}{H_2}\right) + \left(\frac{4}{H_2} - \frac{2}{H_1}\right) = \frac{1}{a} + \frac{2}{b}$$

$$\Rightarrow \frac{3}{H_2} = \frac{b+2a}{ab} \Rightarrow H_2 = \frac{3ab}{2a+b}$$

$$\text{Eq. (iv)} + 2 \times \text{Eq. (iii)} \Rightarrow \left(\frac{2}{H_2} - \frac{1}{H_1}\right) + \left(\frac{4}{H_1} - \frac{2}{H_2}\right) = \frac{1}{b} + \frac{2}{a}$$

$$\Rightarrow \frac{3}{H_1} = \frac{a+2b}{ab} \Rightarrow H_1 = \frac{3ab}{a+2b}$$

$$\therefore \frac{G_1 G_2}{H_1 H_2} = \frac{ab}{\frac{3ab}{a+2b} \times \frac{3ab}{2a+b}} = \frac{(a+2b)(2a+b)}{9ab} \text{ Hence proved.}$$

Ex. 14. If $x > 1, y > 1, z > 1$ are in G.P., then show that $\frac{1}{1 + \log x}, \frac{1}{1 + \log y}, \frac{1}{1 + \log z}$ are in H.P.

(IIT 1999, Manipal 2012)

Sol. x, y, z are in G.P.

$$\Rightarrow y^2 = xz \Rightarrow 2 \log y = \log x + \log z$$

$$\Rightarrow \log x, \log y, \log z \text{ are in A.P.}$$

$$\Rightarrow 1 + \log x, 1 + \log y, 1 + \log z \text{ are in A.P.}$$

(Adding 1 to each term)

$$\Rightarrow \frac{1}{1 + \log x}, \frac{1}{1 + \log y}, \frac{1}{1 + \log z} \text{ are in H.P.}$$

Ex. 15. Find the harmonic mean of the roots of the equation

$$(5 + \sqrt{2})x^2 - (4 + \sqrt{5})x + (8 + 2\sqrt{5}) = 0.$$

(IIT 1999, MPPET 2010, EAMCET 2013)

Sol. Let the roots of the equation be α and β . Then,

$$\text{Sum of roots} = \alpha + \beta = \frac{(4 + \sqrt{5})}{5 + \sqrt{2}}$$

$$\text{Product of roots} = \alpha\beta = \frac{8 + 2\sqrt{5}}{5 + \sqrt{2}}$$

$$\text{Now, Harmonic Mean of the roots, } \alpha \text{ and } \beta = \frac{2\alpha\beta}{\alpha + \beta} = \frac{2 \left(\frac{8 + 2\sqrt{5}}{5 + \sqrt{2}} \right)}{\frac{(4 + \sqrt{5})}{5 + \sqrt{2}}} = \frac{4(4 + \sqrt{5})}{(4 + \sqrt{5})} = 4.$$

PRACTICE SHEET

1. The n th term of the H.P. $4 + 4\frac{2}{7} + 4\frac{8}{13} + 5 + \dots$ is

- (a) $\frac{7}{26-2n}$ (b) $\frac{3}{3n-2}$
 (c) $\frac{60}{16-n}$ (d) None of these

2. The harmonic mean of $\frac{a}{1-ab}$ and $\frac{a}{1+ab}$ is

- (a) a (b) $\frac{1}{1-a^2b^2}$
 (c) $\frac{a}{1-a^2b^2}$ (d) $\frac{a}{\sqrt{1-a^2b^2}}$
 (AMU 2002)

3. The 5th and 11th term of an H.P. are $\frac{1}{45}$ and $\frac{1}{69}$ respectively.

Then, its 16th term will be

- (a) $\frac{1}{77}$ (b) $\frac{1}{81}$ (c) $\frac{1}{85}$ (d) $\frac{1}{89}$
 (Rajasthan PET 2003)

4. If the A.M. and G.M. of two numbers be 27 and 18 respectively, then what is their H.M. equal to?

- (a) 24 (b) 12 (c) 16 (d) 28

5. If the first two terms of an H.P. are $\frac{2}{5}$ and $\frac{12}{13}$ respectively, then the largest term is

- (a) 2nd term (b) 3rd term (c) 4th term (d) 6th term
 (AMU 2007)

6. If a and b are two real numbers such that $0 < a < b$ and the arithmetic mean between a and b is $\frac{4}{3}$ times the harmonic mean between them, then b/a is equal to

- (a) $\frac{2}{\sqrt{3}}$ (b) $\frac{3}{2}$ (c) 3 (d) $\frac{8}{3}$

7. G.M. and H.M. of two numbers are 10 and 8 respectively. The numbers are

- (a) 1, 100 (b) 2, 50 (c) 4, 25 (d) 5, 20
 (WBJEE 2010)

8. Five numbers are in H.P. The middle term is 1 and the ratio of the second and fourth terms is 2 : 1. Then, the sum of the first three terms is

- (a) $11/2$ (b) 5 (c) 2 (d) $14/2$
 (WBJEE 2013)

9. If for two numbers the ratio of their H.M. to G.M. is 20:29, then the numbers are in the ratio

- (a) 3 : 40 (b) 4 : 25 (c) 1 : 22 (d) 2 : 27
 (Type IIT)

10. If $2(y-a)$ is the H.M. between $y-x$ and $y-z$, then $(x-a)$, $(y-a)$, $(z-a)$ are in

- (a) A.P. (b) G.P. (c) H.P. (d) None of these
 (Rajasthan PET 2001)

11. If a, b, c are in H.P., then $\left(\frac{1}{a} + \frac{1}{b} - \frac{1}{c}\right)\left(\frac{1}{b} + \frac{1}{c} - \frac{1}{a}\right)$ is equal to

- (a) $\frac{4}{b^2} - \frac{3}{ac}$ (b) $\frac{3}{b^2} - \frac{4}{ac}$ (c) $\frac{4}{ac} - \frac{3}{b^2}$ (d) $\frac{3}{b^2} + \frac{4}{ac}$
 (Kerala PET)

12. If the l th, m th and n th terms of an H.P. are in H.P. then l, m, n , are in

- (a) H.P. (b) A.P. (c) G.P. (d) None of these

13. If $\log(a+c)$, $\log(c-a)$ and $\log(a-2b+c)$ are in A.P. then

- (a) a, b, c are in A.P. (b) a^2, b^2, c^2 are in A.P.
 (c) a, b, c are in G.P. (d) a, b, c are in H.P.

(DCE 2002)

14. If the sum of the roots of the equation $ax^2 + bx + c = 0$ is equal to sum of their squares, then $\frac{c}{a}, \frac{b}{a}, \frac{c}{b}$ are in

- (a) A.P. (b) G.P. (c) H.P. (d) None of these
 (Punjab CET 2008)

15. If $a^x = b^y = c^z$ and x, y, z are in H.P., then a, b, c are in

- (a) G.P. (b) A.P. (c) H.P. (d) None of these

16. If $x^2 + 9y^2 + 25z^2 = xyz\left(\frac{15}{x} + \frac{5}{y} + \frac{3}{z}\right)$, then x, y, z are in

- (a) G.P. (b) H.P. (c) A.P. (d) None of these
 (DCE 2004)

17. If a, b, c are in A.P. and a^2, b^2, c^2 are in H.P., then which of the following statement can be true?

- (a) $a, b, -\frac{c}{2}$ are in G.P. (b) $a = b = c$
 (c) Any of these (d) None of these

(IIT 2003)

18. The H.M. of two numbers is 4. Their A.M. is A and G.M. is G . If $2A + G^2 = 27$, then A is equal to

- (a) $\frac{9}{2}$ (b) 18 (c) $\frac{27}{2}$ (d) 27

(WBJEE 2011)

19. Let the positive numbers a, b, c, d be in A.P. Then, abc, abd, acd, bcd are

- (a) NOT in A.P./G.P./H.P. (b) In A.P.
 (c) In G.P. (d) In H.P. (IIT 2001)

20. If $a^x = b^y = c^z = d^u$ and a, b, c, d are in GP, then x, y, z, u are in

- (a) A.P. (b) G.P. (c) H.P. (d) None of these
 (VITEEE 2010)

21. If $a, a_1, a_2, \dots, a_{2n}, b$ are in arithmetic progression and $a, g_1, g_2, \dots, g_{2n}, b$ are in geometric progression, and h is the harmonic mean of a and b , then $\frac{a_1 + a_{2n}}{g_1 g_{2n}} + \frac{a_2 + a_{2n-1}}{g_2 g_{2n-1}} + \dots + \frac{a_n + a_{n+1}}{g_n g_{n+1}}$ is equal to
 (a) $2nh$ (b) n/h (c) nh (d) $2n/h$

(DCE 2009)

22. Find the value of n for which $\left(\frac{a^{n+1} + b^{n+1}}{a^n + b^n}\right)$ is the harmonic mean between a and b .
 (a) -2 (b) $-\frac{3}{2}$ (c) -1 (d) $-\frac{1}{2}$

23. If $\frac{1}{b-a} + \frac{1}{b-c} = \frac{1}{a} + \frac{1}{c}$ and $b \neq (a+c)$, then a, b, c are in
 (a) A.P (b) G.P (c) H.P (d) None of these
24. If a, b, c are in H.P. and half the middle term be subtracted from the three terms, then the resulting series will be in
 (a) A.P. (b) G.P (c) H.P (d) None of these
25. If p, q, r are in H.P and the $(p+1)$ th, $(q+1)$ th and $(r+1)$ th terms of an A.P. are in G.P., then the ratio of the first term to the common difference of the A.P. is equal to
 (a) $\frac{-q}{2}$ (b) $\frac{-pr}{q}$ (c) $\frac{-pr}{q^2}$ (d) $\frac{-2q}{pr}$

ANSWERS

1. (c) 2. (a) 3. (d) 4. (b) 5. (a) 6. (c) 7. (d) 8. (a) 9. (b) 10. (b)
 11. (c) 12. (b) 13. (d) 14. (c) 15. (a) 16. (b) 17. (c) 18. (a) 19. (d) 20. (c)
 21. (d) 22. (c) 23. (c) 24. (b) 25. (a)

HINTS AND SOLUTIONS

1. $4, 4\frac{2}{7}, 4\frac{8}{13}, 5, \dots$ are in H.P.
 $\Rightarrow \frac{1}{4}, \frac{7}{30}, \frac{13}{60}, \frac{1}{5}, \dots$ are in A.P.
 Now, here $a = \frac{1}{4}, d = \frac{7}{30} - \frac{1}{4} = -\frac{1}{60}$
 $\therefore T_n = a + (n-1)d = \frac{1}{4} + (n-1)\left(-\frac{1}{60}\right)$
 $= \frac{1}{4} - \frac{n}{60} + \frac{1}{60} = \frac{15+1-n}{60} = \frac{16-n}{60}$
 \therefore n th term of the H.P. = $\frac{60}{16-n}$.
2. \therefore Harmonic mean of two quantities a and b is $\frac{2ab}{a+b}$
 \therefore Reqd. H.M. = $\frac{2 \times \left(\frac{a}{1-ab}\right) \times \left(\frac{a}{1+ab}\right)}{\frac{a}{1-ab} + \frac{a}{1+ab}}$
 $= \frac{2a^2}{\frac{(1-a^2b^2)}{1-a^2b^2}} = \frac{2a^2}{1-a^2b^2} = \frac{2a^2}{a} = a$.
3. Given, 5th term of an H.P. = $\frac{1}{45}$
 \Rightarrow 5th term of corresponding A.P. = 45
 11th term of an H.P. = $\frac{1}{69}$

- \Rightarrow 11th term of corresponding A.P. = 69
 Let a and d be the first term and common difference of this A.P.
 So $a + 4d = 45$... (i)
 $a + 10d = 69$... (ii)
 (ii) - (i) $\Rightarrow 6d = 24 \Rightarrow d = 4$
 From (i) $a + 16 = 45 \Rightarrow a = 29$.
 $\therefore T_{16} = a + 15d = 29 + 15 \times 4 = 29 + 69 = 89$
 \therefore 16th term of corresponding H.P. = $\frac{1}{89}$.
4. $\therefore (G.M)^2 = (A.M) \cdot (H.M)$
 $\therefore (18)^2 = 27 \times H.M. \Rightarrow H.M. = \frac{18 \times 18}{27} = 12$.
5. First term of H.P = $\frac{2}{5} \Rightarrow$ First term of A.P = $\frac{5}{2} = \frac{30}{12}$
 Second term of H.P = $\frac{12}{13} \Rightarrow$ Second term of A.P = $\frac{13}{12}$
 \therefore The first and second terms of the corresponding A.P. are $\frac{30}{12}$ and $\frac{13}{12}$ respectively. The common difference
 $d = \frac{13}{12} - \frac{30}{12} = -\frac{17}{12}$
 \therefore The corresponding A.P. is $\frac{30}{12}, \frac{13}{12}, -\frac{4}{12}, -\frac{21}{12}, \dots$
 \therefore The corresponding H.P is $\frac{12}{30}, \frac{12}{13}, -\frac{12}{4}, -\frac{12}{21}, \dots$
 All the terms after the second term are negative, so the largest term is out of the first two terms, i.e., between $\frac{12}{30}$ and $\frac{12}{13}$.

$\therefore \frac{12}{13} > \frac{12}{30}$, so, $\frac{12}{13}$ i.e., second term is the largest term of the H.P.

6. Arithmetic mean between a and b is $A = \frac{a+b}{2}$

Harmonic mean between a and b is $H = \frac{2ab}{a+b}$

$$\text{Given, } A = \frac{4}{3}H \Rightarrow \frac{a+b}{2} = \frac{4}{3} \left(\frac{2ab}{a+b} \right)$$

$$\Rightarrow 3(a+b)^2 = 16ab \Rightarrow 3b^2 - 10ab + 3a^2 = 0$$

$$\Rightarrow 3 \left(\frac{b}{a} \right)^2 - 10 \left(\frac{b}{a} \right) + 3 = 0 \quad (\text{Dividing throughout by } a^2)$$

$$\Rightarrow \left(\frac{3b}{a} - 1 \right) \left(\frac{b}{a} - 3 \right) = 0$$

$$\Rightarrow \frac{b}{a} = \frac{1}{3} \text{ or } \frac{b}{a} = 3 \quad \left[\because 0 < a < b \Rightarrow \frac{b}{a} \neq \frac{1}{3} \right]$$

$$\Rightarrow \frac{b}{a} = 3.$$

7. Let the two required numbers be a and b .

$$\text{Then, G.M} = \sqrt{ab} = 10 \Rightarrow ab = 100 \quad \dots(i)$$

$$\text{H.M.} = \frac{2ab}{a+b} = 8 \Rightarrow a+b = \frac{ab}{4} = \frac{100}{4} = 25 \quad \dots(ii)$$

$$\therefore (a-b)^2 = (a+b)^2 - 4ab \\ = 625 - 400 = 225$$

$$\Rightarrow a-b = \pm 15 \quad \dots(iii)$$

Solving (i) and (ii), we get $a = 20, b = 5$ or $a = 5, b = 20$

\therefore The required numbers are **20** and **5**.

8. Let the terms of the corresponding A.P. be $a-2d, a-d, a, a+d, a+2d$.

$$\text{Given, } a = 1 \text{ and } \frac{a-d}{a+d} = \frac{1}{2}$$

$$\Rightarrow 2a - 2d = a + d$$

$$\Rightarrow a = 3d \Rightarrow d = \frac{1}{3}a = \frac{1}{3}. \quad (\because a = 1)$$

$$\therefore \text{ First three terms of A.P are } 1 - \frac{2}{3}, 1 - \frac{1}{3}, 1, \text{ i.e., } \frac{1}{3}, \frac{2}{3}, 1$$

$$\Rightarrow \text{ First three terms of corresponding H.P are } 3, \frac{3}{2}, 1$$

$$\therefore \text{ Required sum} = 3 + \frac{3}{2} + 1 = 5\frac{1}{2} = \frac{11}{2}.$$

9. Let the two numbers be a and b .

$$\text{Given, } \frac{\text{H.M}}{\text{G.M}} = \frac{20}{29} \Rightarrow \frac{2ab/a+b}{\sqrt{ab}} = \frac{20}{29}$$

$$\Rightarrow \frac{2\sqrt{ab}}{a+b} = \frac{20}{29} \Rightarrow 58\sqrt{ab} = 20(a+b)$$

$$\Rightarrow 20a - 58\sqrt{ab} + 20b = 0$$

$$\Rightarrow 20\frac{a}{b} - 58\sqrt{\frac{a}{b}} + 20 = 0 \quad (\text{Dividing all terms by } b)$$

$$\Rightarrow 20x^2 - 58x + 20 = 0 \quad (\text{where } x = \sqrt{\frac{a}{b}})$$

$$\Rightarrow 20x^2 - 50x - 8x + 20 = 0$$

$$\Rightarrow 10x(2x-5) - 4(2x-5) = 0$$

$$\Rightarrow (10x-4)(2x-5) = 0$$

$$\Rightarrow x = \frac{2}{5} \text{ or } \frac{5}{2}$$

$$\Rightarrow \sqrt{\frac{a}{b}} = \frac{2}{5} \text{ or } \frac{5}{2}$$

$$\Rightarrow \frac{a}{b} = \frac{4}{25} \text{ or } \frac{25}{4}$$

Thus from the given options, the two numbers are in the ratio **4 : 25**.

10. Given, $2(y-a)$ is the H.M. between $(y-x)$ and $(y-z)$

$$\Rightarrow (y-x), 2(y-a), (y-z) \text{ are in H.P.}$$

$$\Rightarrow \frac{1}{y-x}, \frac{1}{2(y-a)}, \frac{1}{y-z} \text{ are in A.P.}$$

$$\Rightarrow \frac{1}{2(y-a)} - \frac{1}{(y-x)} = \frac{1}{(y-z)} - \frac{1}{2(y-a)}$$

$$\Rightarrow \frac{y-x-2y+2a}{2(y-a)(y-x)} = \frac{2y-2a-y+z}{(y-z)2(y-a)}$$

$$\Rightarrow \frac{-x-y+2a}{(y-x)} = \frac{y+z-2a}{(y-z)}$$

$$\Rightarrow \frac{x+y-2a}{x-y} = \frac{y+z-2a}{y-z}$$

$$\Rightarrow \frac{(x-a)+(y-a)}{(x-a)-(y-a)} = \frac{(y-a)+(z-a)}{(y-a)-(z-a)}$$

Now applying componendo and dividendo, we have

$$\left(\text{By comp. and div., } \frac{x}{y} = \frac{a}{b} \Rightarrow \frac{x+y}{x-y} = \frac{a+b}{a-b} \right)$$

$$\Rightarrow \frac{2(x-a)}{2(y-a)} = \frac{2(y-a)}{2(z-a)}$$

$$\Rightarrow (x-a)(z-a) = (y-a)^2$$

$$\Rightarrow (x-a), (y-a), (z-a) \text{ are in G.P.}$$

11. If a, b, c are in H.P, then $b = \frac{2ac}{a+c}$

$$\therefore \left(\frac{1}{a} + \frac{1}{b} - \frac{1}{c} \right) \left(\frac{1}{b} + \frac{1}{c} - \frac{1}{a} \right)$$

$$= \left(\frac{1}{a} + \frac{a+c}{2ac} - \frac{1}{c} \right) \left(\frac{a+c}{2ac} + \frac{1}{c} - \frac{1}{a} \right)$$

$$= \left(\frac{2c+a+c-2a}{2ac} \right) \left(\frac{a+c+2a-2c}{2ac} \right)$$

$$= \left(\frac{3c-a}{2ac} \right) \left(\frac{3a-c}{2ac} \right) = \frac{10ac - 3a^2 - 3c^2}{4a^2c^2}$$

$$= \frac{16ac - (3a^2 + 3c^2 + 6ac)}{4a^2c^2} = \frac{16ac - 3(a+c)^2}{4a^2c^2}$$

$$= \frac{4}{ac} - 3\left(\frac{a+c}{2ac}\right)^2 = \frac{4}{ac} - \frac{3}{b^2}$$

12. Let the l th, m th and n th terms of the corresponding A.P. be

$$\left. \begin{aligned} T_l &= a + (l-1)d \\ T_m &= a + (m-1)d \\ T_n &= a + (n-1)d \end{aligned} \right\} \Rightarrow \begin{aligned} 2T_m &= T_l + T_n \\ 2[a + (m-1)d] &= a + (l-1)d + a + (n-1)d \\ (2m-2)d &= (l+n-2)d \\ 2m &= l+n \\ l, m, n &\text{ are in A.P.} \end{aligned}$$

13. $\log(a+c)$, $\log(c-a)$ and $\log(a-2b+c)$ are in A.P.

$$\begin{aligned} \Rightarrow 2 \log(c-a) &= \log(a+c) + \log(a-2b+c) \\ \Rightarrow \log(c-a)^2 &= \log[(a+c)(a-2b+c)] \\ \Rightarrow c^2 + a^2 - 2ac &= a^2 + ca - 2ba - 2bc + ac + c^2 \\ \Rightarrow 2ab + 2bc &= 4ac \Rightarrow b(a+c) = 2ac \\ \Rightarrow b &= \frac{2ac}{a+c} \Rightarrow a, b, c \text{ are in H.P.} \end{aligned}$$

14. Let α, β be the roots of the equation $ax^2 + bx + c = 0$.

Then, $\alpha + \beta = -\frac{b}{a}, \alpha\beta = \frac{c}{a}$

Also, given $\alpha + \beta = \alpha^2 + \beta^2 \Rightarrow \alpha + \beta = (\alpha + \beta)^2 - 2\alpha\beta$

$$\Rightarrow -\frac{b}{a} = \left(-\frac{b}{a}\right)^2 - \frac{2c}{a}$$

$$\Rightarrow -ba = b^2 - 2ac \Rightarrow b^2 + ab = 2ac$$

$$\Rightarrow b(b+a) = 2ac \Rightarrow \frac{b}{c} + \frac{a}{c} = \frac{2a}{b}$$

$\Rightarrow \frac{c}{a}, \frac{a}{b}, \frac{b}{c}$ are in A.P.

15. x, y, z are in H.P $\Rightarrow y = \frac{2xz}{x+z}$... (i)

Given, $a^x = b^y = c^z = k$ (say)

Then $x \log a = \log k, y \log b = \log k, z \log c = \log k$

$$\Rightarrow x = \frac{\log k}{\log a}, y = \frac{\log k}{\log b}, z = \frac{\log k}{\log c}$$

\therefore Putting these values of x, y, z in (i), we get

$$\frac{\log k}{\log b} = \frac{2 \left(\frac{\log k}{\log a} \right) \left(\frac{\log k}{\log c} \right)}{\frac{\log k}{\log a} + \frac{\log k}{\log c}}$$

$$\Rightarrow \frac{\log k}{\log b} = \frac{2(\log k)^2}{\log k \log c + \log a \log k}$$

$$\Rightarrow \frac{(\log k)^2}{\log b} = \frac{2(\log k)^2}{\log c + \log a}$$

$$\Rightarrow 2 \log b = \log c + \log a \Rightarrow \log b^2 = \log(ac)$$

$$\Rightarrow b^2 = ac \Rightarrow a, b, c \text{ are in G.P.}$$

16. Given, $x^2 + 9y^2 + 25z^2 = xyz \left(\frac{15}{x} + \frac{5}{y} + \frac{3}{z} \right)$

$$\begin{aligned} \Rightarrow x^2 + 9y^2 + 25z^2 - 15yz - 5xz - 3xy &= 0 \\ \Rightarrow 2x^2 + 18y^2 + 50z^2 - 30yz - 10xz - 6xy &= 0 \\ \Rightarrow (x-3y)^2 + (3y-5z)^2 + (x-5z)^2 &= 0 \\ \Rightarrow x=3y, 3y=5z, 5z=x \Rightarrow x=3y=5z \\ \Rightarrow \frac{x}{1} = \frac{y}{\frac{1}{3}} = \frac{z}{\frac{1}{5}} \Rightarrow \frac{x}{15} = \frac{y}{5} = \frac{z}{3} \end{aligned}$$

$$\Rightarrow x, y, z \text{ are in H.P.} \left(\begin{aligned} \because y &= \frac{2xz}{x+z} \\ y &= \frac{2 \times 15 \times 3}{15+3} = 5 \end{aligned} \right)$$

17. a, b, c are in A.P. $\Rightarrow 2b = a+c$... (i)

$$a^2, b^2, c^2 \text{ are in H.P.} \Rightarrow b^2 = \frac{2a^2c^2}{a^2+c^2} \dots (ii)$$

From eqn (ii)

$$\begin{aligned} b^2(a^2+c^2) &= 2a^2c^2 \Rightarrow b^2\{(a+c)^2-2ac\} = 2a^2c^2 \\ \Rightarrow b^2\{4b^2-2ac\} &= 2a^2c^2 \quad (\text{From (i) } 2b = a+c) \\ \Rightarrow 2b^4 - b^2ac - a^2c^2 &= 0 \dots (iii) \\ \Rightarrow (b^2-ac)(2b^2+ac) &= 0 \\ \Rightarrow b^2-ac=0 \text{ or } 2b^2+ac &= 0 \\ \Rightarrow \left(\frac{a+c}{2}\right)^2 - ac = 0 \quad b^2 = -\frac{ac}{2} \\ \Rightarrow (a-c)^2 = 0 \quad a, b, -\frac{c}{2} &\text{ are in G.P.} \\ \Rightarrow 2b = 2c \quad (\text{From (i)}) \\ \Rightarrow b &= c. \end{aligned}$$

18. Let the two numbers be a and b . Then,

$$\text{H.M} = \frac{2ab}{a+b} = 4 \Rightarrow a+b = \frac{ab}{2} \dots (i)$$

Also given, $A = \frac{a+b}{2}, G = \sqrt{ab}$

$$\begin{aligned} \therefore 2A + G^2 = 27 \Rightarrow 2 \times \frac{a+b}{2} + ab &= 27 \\ \Rightarrow a+b+ab = 27 \Rightarrow (a+b) + 2(a+b) &= 27 \\ \Rightarrow 3(a+b) = 27 \Rightarrow a+b=9 \Rightarrow \frac{a+b}{2} = \frac{9}{2} = A. \end{aligned}$$

19. If a, b, c, d are in A.P.

$$\begin{aligned} \Rightarrow d, c, b, a &\text{ are in A.P.} \\ \Rightarrow \frac{d}{abcd}, \frac{c}{abcd}, \frac{b}{abcd}, \frac{a}{abcd} &\text{ are in A.P.} \\ &(\text{Dividing all terms by } abcd) \\ \Rightarrow \frac{1}{abc}, \frac{1}{abd}, \frac{1}{acd}, \frac{1}{bcd} &\text{ are in A.P.} \\ \Rightarrow abc, abd, acd, bcd &\text{ are in H.P.} \end{aligned}$$

20. Given, $a^x = b^y = c^z = d^u = k$ (say)

Then, $a = k^{1/x}, b = k^{1/y}, c = k^{1/z}, d = k^{1/u}$... (i)

Since a, b, c, d are in G.P.

$$\frac{b}{a} = \frac{c}{b} = \frac{d}{c}$$

$$\Rightarrow \frac{k^{1/y}}{k^{1/x}} = \frac{k^{1/z}}{k^{1/y}} = \frac{k^{1/u}}{k^{1/z}} \quad (\text{Using (i)})$$

$$\Rightarrow k^{1/y - 1/x} = k^{1/z - 1/y} = k^{1/u - 1/z}$$

$$\Rightarrow \frac{1}{y} - \frac{1}{x} = \frac{1}{z} - \frac{1}{y} = \frac{1}{u} - \frac{1}{z}$$

$$\Rightarrow \frac{1}{x}, \frac{1}{y}, \frac{1}{z}, \frac{1}{u} \text{ are in A.P.} \Rightarrow x, y, z, u \text{ are in H.P.}$$

21. Given, $a, a_1, a_2, a_3, \dots, a_{2n}, b$ are in A.P.

$$\Rightarrow a_1 + a_{2n} = a_2 + a_{2n-1} = \dots = a + b$$

$$(\because a_1 - a = b - a_{2n}, a_2 - a = b - a_{2n-1})$$

Also, $a, g_1, g_2, \dots, g_{2n}, b$ are in G.P.

$$\Rightarrow g_1 g_{2n} = g_2 g_{2n-1} = \dots = ab$$

$$\therefore \frac{a_1 + a_{2n}}{g_1 g_{2n}} + \frac{a_2 + a_{2n-1}}{g_2 g_{2n-1}} + \dots + \frac{a_n + a_{n+1}}{g_n g_{n+1}}$$

$$= \frac{a+b}{ab} + \frac{a+b}{ab} + \dots + \frac{a+b}{ab} \quad (n \text{ times})$$

$$= \frac{n(a+b)}{ab} = \frac{2n}{h}, \text{ where,}$$

[Given, $h = \frac{2ab}{a+b}$ is the harmonic mean between a and b]

22. If $\frac{a^{n+1} + b^{n+1}}{a^n + b^n}$ is the harmonic mean between a and b , then

$$\frac{a^{n+1} + b^{n+1}}{a^n + b^n} = \frac{2ab}{a+b}$$

$$\Rightarrow a^{n+1} \cdot a + a^{n+1} b + b^{n+1} a + b^{n+1} \cdot b = 2ab \cdot a^n + 2ab \cdot b^n$$

$$\Rightarrow a^{n+2} + a^{n+1} b + b^{n+1} a + b^{n+2} = 2a^{n+1} b + 2ab^{n+1}$$

$$\Rightarrow a^{n+2} + b^{n+2} = a^{n+1} b + ab^{n+1}$$

$$\Rightarrow a^{n+2} - ba^{n+1} = ab^{n+1} - b^{n+2}$$

$$\Rightarrow a^{n+1}(a-b) = b^{n+1}(a-b)$$

$$\Rightarrow a^{n+1} = b^{n+1} \Rightarrow \left(\frac{a}{b}\right)^{n+1} = 1 \Rightarrow \left(\frac{a}{b}\right)^{n+1} = \left(\frac{a}{b}\right)^0$$

$$\Rightarrow n+1=0 \Rightarrow n=-1.$$

23. $\frac{1}{b-a} + \frac{1}{b-c} = \frac{1}{a} + \frac{1}{c} \Rightarrow \frac{b-c+b-a}{(b-a)(b-c)} = \frac{c+a}{ac}$

$$\Rightarrow \frac{2b-(c+a)}{b^2-b(a+c)+ac} = \frac{a+c}{ac}$$

$$\Rightarrow 2bac - ac(a+c) = b^2(a+c) - b(a+c)^2 + ac(a+c)$$

$$\Rightarrow b^2(a+c) - b(a+c)^2 + 2ac(a+c) - 2abc = 0$$

$$\Rightarrow b(a+c)(b-(a+c)) + 2ac(a+c-b) = 0$$

$$\Rightarrow b(a+c)(b-(a+c)) - 2ac(b-(a+c)) = 0$$

$$\Rightarrow [b(a+c) - 2ac](b-(a+c)) = 0$$

$$\Rightarrow b(a+c) - 2ac = 0 (\because b \neq a+c \Rightarrow b-(a+c) \neq 0)$$

$$\Rightarrow b = \frac{2ac}{a+c} \Rightarrow a, b, c \text{ are in H.P.}$$

24. Let the three numbers in H.P. be a, b, c .

$$\therefore b = \frac{2ac}{a+c}$$

The new numbers after subtracting half the middle term from each term are:

$$a - \frac{b}{2} = a - \frac{ac}{a+c} = \frac{a^2}{a+c}$$

$$b - \frac{b}{2} = \frac{b}{2} = \frac{ac}{a+c}$$

$$c - \frac{b}{2} = c - \frac{ac}{a+c} = \frac{c^2}{a+c}$$

$$\text{Now, } \left(a - \frac{b}{2}\right) \left(c - \frac{b}{2}\right) = \left(\frac{a^2}{a+c}\right) \left(\frac{c^2}{a+c}\right)$$

$$= \frac{a^2 c^2}{(a+c)^2} = \left(\frac{b}{2}\right)^2$$

$$\Rightarrow \left(a - \frac{b}{2}\right), \frac{b}{2}, \left(c - \frac{b}{2}\right) \text{ are in G.P.}$$

25. p, q, r are in H.P. $\Rightarrow q = \frac{2pr}{p+r} \dots(i)$

Let a and d be the first term and common difference respectively of the A.P.

$$\therefore T_{p+1} = a + pd$$

$$T_{q+1} = a + qd$$

$$T_{r+1} = a + rd$$

Given, $T_{p+1}, T_{q+1}, T_{r+1}$ are in G.P.

$$\Rightarrow (T_{q+1})^2 = T_{p+1} \cdot T_{r+1}$$

$$(a + qd)^2 = (a + pd)(a + rd)$$

$$\Rightarrow a^2 + 2aqd + q^2 d^2 = a^2 + ad(p+r) + prd^2$$

$$\Rightarrow ad(2q-p-r) = d^2(pr-q^2)$$

$$\Rightarrow \frac{a}{d} = \frac{pr-q^2}{2q-p-r} = \frac{\frac{1}{2}(p+r)q - q^2}{2q-(p+r)} \quad (\text{Using (i)})$$

$$= \frac{-\frac{q}{2}[2q-(p+r)]}{2q-(p+r)} = -\frac{q}{2}$$

ARITHMETICO-GEOMETRIC SERIES

KEY FACTS

1. A series in which each term is the product of corresponding terms of an A.P. and a G.P. is called an **Arithmetico-Geometric series**.

The **general** or **standard** form of such a series is

$a + (a + d)r + (a + 2d)r^2 + (a + 3d)r^3 + \dots + \{a + (n - 1)d\}r^{n-1} + \dots$ where each term is formed by multiplying the corresponding terms of the two series.

A.P. : $a + (a + d) + (a + 2d) + \dots + (a + (n - 1)d) + \dots$ and

G.P. : $1 + r + r^2 + \dots + r^{n-1} + \dots$

2. n th term of an Arithmetico-Geometric series is

$$T_n = \{a + (n - 1)d\} r^{n-1}.$$

3. **Sum of n terms of an Arithmetico-Geometric Series**

Let S_n be the sum of the n terms of the series $a + (a + d)r + (a + 2d)r^2 + \dots + (a + (n - 1)d)r^{n-1}$

$$\text{Then, } S_n = a + (a + d)r + (a + 2d)r^2 + \dots + (a + (n - 1)d)r^{n-1} \quad \dots(i)$$

$$rS_n = ar + (a + d)r^2 + \dots + (a + (n - 2)d)r^{n-1} + (a + (n - 1)d)r^n \quad \dots(ii)$$

Subtracting eqn (ii) from eqn (i), we get

$$(1 - r)S_n = (a + dr + dr^2 + \dots + dr^{n-1}) - (a + (n - 1)d)r^n$$

$$= a + \frac{dr(1 - r^{n-1})}{(1 - r)} - (a + (n - 1)d)r^n$$

$$\Rightarrow S_n = \frac{a}{1 - r} + \frac{dr(1 - r^{n-1})}{(1 - r)^2} - \frac{(a + (n - 1)d)r^n}{(1 - r)}.$$

4. **Sum of an infinite Arithmetico-Geometric Series**

Let the infinite Arithmetico-Geometric series be

$$a + (a + d)r + (a + 2d)r^2 + \dots + (a + (n - 1)d)r^{n-1} + \dots \infty$$

If $|r| < 1$, then $r^n \rightarrow 0$ and $r^{n-1} \rightarrow 0$ as $n \rightarrow \infty$, then

$$S_\infty = \frac{a}{1 - r} + \frac{dr}{(1 - r)^2}.$$

SOLVED EXAMPLES

Ex. 1. Find the n th term of the given arithmetico-geometric series:

$$(i) \frac{1}{3} + \frac{3}{9} + \frac{5}{27} + \frac{7}{81} + \dots \quad (ii) 1 - 2x + 3x^2 - 4x^3 + \dots$$

Sol. (i) The A.P. and G.P. corresponding to the given series

$$\frac{1}{3} + \frac{3}{9} + \frac{5}{27} + \frac{7}{81} + \dots \text{ are respectively}$$

$$1, 3, 5, 7, \dots \text{ and } \frac{1}{3}, \frac{1}{9}, \frac{1}{27}, \frac{1}{81}, \dots$$

$$n\text{th term of A.P.} = (1 + (n - 1)2) = (2n - 1)$$

$$n\text{th term of G.P.} = \frac{1}{3} \cdot \left(\frac{1}{3}\right)^{n-1} = \left(\frac{1}{3}\right)^n$$

$$\therefore n\text{th term of the given series} = (2n - 1) \left(\frac{1}{3}\right)^n.$$

(ii) The given arithmetico-geometric series is

$$1 - 2x + 3x^2 - 4x^3 + \dots$$

where corresponding A.P. and G.P. are respectively 1, 2, 3, 4, and 1, $-x$, $(-x)^2$, $(-x)^3$,

$$n\text{th term of A.P.} = (1 + (n - 1) 1) = n$$

$$n\text{th term of G.P.} = 1 \cdot (-x)^{n-1} = (-1)^{n-1} (x)^{n-1}.$$

$$\therefore n\text{th term of the given series} = n \cdot (-1)^{n-1} x^{n-1} = (-1)^{n-1} nx^{n-1}.$$

Ex. 2. Find the sum to n terms of the series $1 + \frac{3}{2} + \frac{5}{4} + \frac{7}{8} + \dots$

Sol. $1 + \frac{3}{2} + \frac{5}{4} + \frac{7}{8} + \dots$ is an arithmetico-geometric series with corresponding A.P. and G.P. as:

$$\text{A.P. : } 1 + 3 + 5 + 7 + \dots$$

$$\text{G.P. : } 1 + \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots$$

$$n\text{th term of A.P.} = (1 + (n - 1) 2) = (2n - 1)$$

$$n\text{th term of G.P.} = 1 \cdot \left(\frac{1}{2}\right)^{n-1} = \frac{1}{2^{n-1}}$$

$$\therefore n\text{th term of the given A.G.P.} = \frac{2n-1}{2^{n-1}}$$

$$\therefore S_n = 1 + \frac{3}{2} + \frac{5}{4} + \frac{7}{8} + \dots + \frac{2n-1}{2^{n-1}}$$

$$\Rightarrow \frac{1}{2} S_n = \frac{1}{2} + \frac{3}{4} + \frac{5}{8} + \dots + \frac{2n-3}{2^{n-1}} + \frac{2n-1}{2^n}$$

On subtraction, we get

$$\Rightarrow \left(1 - \frac{1}{2}\right) S_n = 1 + \frac{2}{2} + \frac{2}{4} + \frac{2}{8} + \dots + \frac{2}{2^{n-1}} - \frac{2n-1}{2^n}$$

$$\Rightarrow \frac{1}{2} S_n = 1 + 2 \left\{ \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots + \frac{1}{2^{n-1}} \right\} - \frac{2n-1}{2^n}$$

$$\Rightarrow \frac{1}{2} S_n = 1 + 2 \left\{ \frac{\left(\frac{1}{2}\right) \left(1 - \left(\frac{1}{2}\right)^n\right)}{1 - \frac{1}{2}} \right\} - \frac{2n-1}{2^n} = 1 + 2 - \frac{4}{2^n} - \frac{2n-1}{2^n} = 3 - \frac{2n+3}{2^n}$$

$$\therefore S_n = 6 - \frac{2n+3}{2^{n-1}}.$$

Ex. 3. Find the sum of the series $1 + \frac{4}{5} + \frac{7}{5^2} + \frac{10}{5^3} + \dots \infty$

Sol. Let $S_\infty = 1 + \frac{4}{5} + \frac{7}{5^2} + \frac{10}{5^3} + \dots \infty$... (i)

$$\frac{1}{5} S_\infty = \frac{1}{5} + \frac{4}{5^2} + \frac{7}{5^3} + \dots \infty$$
 ... (ii)

Subtracting eqn (ii) from eqn (i), we get

$$\left(1 - \frac{1}{5}\right) S_\infty = 1 + \frac{3}{5} + \frac{3}{5^2} + \frac{3}{5^3} + \dots \infty$$

$$\begin{aligned} \frac{4}{5}S_{\infty} &= 1 + 3\left(\frac{1}{5} + \frac{1}{5^2} + \frac{1}{5^3} + \dots \infty\right) \\ &= 1 + 3\left(\frac{\frac{1}{5}}{1 - \frac{1}{5}}\right) \quad (\because \text{Sum of infinite G.P} = \frac{a}{1-r}) = 1 + \frac{3/5}{4/5} = 1 + \frac{3}{4} = \frac{7}{4} \\ \therefore S_{\infty} &= \frac{7}{4} \times \frac{5}{4} = \frac{35}{16}. \end{aligned}$$

Ex. 4. If the sum to infinity of the series $3 + 5r + 7r^2 + \dots \infty$ is $\frac{44}{9}$, find the value of r .

Sol. $3 + 5r + 7r^2 + \dots \infty$ is an infinite arithmetico-geometric series, where

$$a = 3, d = 2, \text{ common ratio } (r) = r.$$

Sum to infinity of an A.G.P., with first term of A.P. as a , common difference d and common ratio r is

$$\begin{aligned} S_{\infty} &= \frac{a}{1-r} + \frac{dr}{(1-r)^2} \\ \therefore \frac{44}{9} &= \frac{3}{1-r} + \frac{2r}{(1-r)^2} \Rightarrow \frac{44}{9} = \frac{3(1-r) + 2r}{(1-r)^2} \\ \Rightarrow 44(1-r)^2 &= 9(3-r) \Rightarrow 44(1-2r+r^2) = 27-9r \\ \Rightarrow 44-88r+44r^2 &= 27-9r \Rightarrow 44r^2-79r+17=0 \\ \Rightarrow (4r-1)(11r-17) &= 0 \Rightarrow r = \frac{1}{4} \text{ or } \frac{17}{11} \\ r \neq \frac{17}{11} &\text{ as it is not possible to find the sum of an infinite G.P. with } |r| > 1. \text{ So } r = \frac{1}{4}. \end{aligned}$$

Ex. 5. Show that $2^{\frac{1}{4}} \times 4^{\frac{1}{8}} \times 8^{\frac{1}{16}} \times 16^{\frac{1}{32}} \times \dots \infty = 2$

Sol. Let $x = 2^{\frac{1}{4}} \times 4^{\frac{1}{8}} \times 8^{\frac{1}{16}} \times 16^{\frac{1}{32}} \times \dots \infty$

$$\begin{aligned} \therefore \log x &= \frac{1}{4} \log 2 + \frac{1}{8} \log 4 + \frac{1}{16} \log 8 + \frac{1}{32} \log 16 + \dots \infty \\ &= \frac{1}{4} \log 2 + \frac{1}{8} \log 2^2 + \frac{1}{16} \log 2^3 + \frac{1}{32} \log 2^4 + \dots \infty \\ &= \frac{1}{4} \log 2 + \frac{2}{8} \log 2 + \frac{3}{16} \log 2 + \frac{4}{32} \log 2 + \dots \infty \\ &= \left(\frac{1}{4} + \frac{2}{8} + \frac{3}{16} + \frac{4}{32} + \dots \infty\right) \log 2 \quad \dots(i) \end{aligned}$$

Now, $\frac{1}{4} + \frac{2}{8} + \frac{3}{16} + \frac{4}{32} + \dots \infty$ is an Arithmetico-Geometric series.

Let $S = \frac{1}{4} + \frac{2}{8} + \frac{3}{16} + \frac{4}{32} + \dots \infty \quad \dots(ii)$

$$\frac{1}{2}S = \frac{1}{8} + \frac{2}{16} + \frac{3}{32} + \dots \infty \quad \dots(iii)$$

On subtracting eqn (iii) from eqn (ii), we get

$$\frac{1}{2}S = \frac{1}{4} + \frac{1}{8} + \frac{1}{16} + \frac{1}{32} + \dots \infty = \frac{\frac{1}{4}}{1 - \frac{1}{2}} = \frac{\frac{1}{4}}{\frac{1}{2}} = \frac{1}{2} \Rightarrow S = 1$$

$$\begin{aligned} \therefore \log x &= 1 \times \log 2 \\ \Rightarrow \log x &= \log 2 \Rightarrow x = 2. \end{aligned} \quad \text{(From (i))}$$

PRACTICE SHEET

1. The sum to n terms of the series $1 + \frac{4}{5} + \frac{7}{5^2} + \frac{10}{5^3} + \dots$ is

$$(a) \left(\frac{25}{14} - \frac{7n+10}{14 \times 5^{n-1}} \right) \quad (b) \left(\frac{17}{12} - \frac{4n+7}{12 \times 5^{n-1}} \right)$$

$$(c) \left(\frac{35}{16} - \frac{12n+7}{16 \times 5^{n-1}} \right) \quad (d) \left(\frac{15}{11} - \frac{10n+2}{11 \times 5^{n-1}} \right)$$

2. Sum the series: $1 + 2.2 + 3.2^2 + \dots + 100.2^{99}$

$$(a) 99.2^{100} \quad (b) 100.2^{100}$$

$$(c) 99.2^{100} + 1 \quad (d) 1000.2^{100}$$

(Kerala PET 2001)

3. The sum to infinity of the series $1 + \frac{2}{3} + \frac{6}{3^2} + \frac{10}{3^3} + \frac{14}{3^4} + \dots$ is

$$(a) 6 \quad (b) 2 \quad (c) 3 \quad (d) 4$$

(AIEEE 2009)

4. Given $\cot \theta = 2\sqrt{2}$, the sum of the infinite series $1 + 2(1 - \sin \theta) + 3(1 - \sin \theta)^2 + 4(1 - \sin \theta)^3 + \dots$ is

$$(a) 6\sqrt{2} \quad (b) 8 \quad (c) 9 \quad (d) 8\sqrt{2}$$

(AMU 2006)

5. If $4 + \frac{4+d}{5} + \frac{4+2d}{5^2} + \dots \infty = 10$, then d is equal to

$$(a) 5 \quad (b) 8 \quad (c) 10 \quad (d) 16$$

(DCE 2007)

ANSWERS

1. (c) 2. (c) 3. (b) 4. (c) 5. (d)

HINTS AND SOLUTIONS

1. $1 + \frac{4}{5} + \frac{7}{5^2} + \frac{10}{5^3} + \dots$ is an A.G.P. with

$$\text{A.P. : } 1 + 4 + 7 + 10 + \dots$$

$$\text{G.P. : } 1 + \frac{1}{5} + \frac{1}{5^2} + \frac{1}{5^3} + \dots$$

$$\text{nth term of A.P.} = 1 + 3(n-1) = 3n-2$$

$$\text{nth term of G.P.} = 1 \times \left(\frac{1}{5}\right)^{n-1} = \frac{1}{5^{n-1}}$$

$$\therefore \text{nth term of the A.G.P.} = \frac{3n-2}{5^{n-1}}$$

$$(n-1)\text{th term of the A.G.P.} = \frac{3(n-1)-2}{5^{n-2}} = \frac{3n-5}{5^{n-2}}$$

$$\text{Now let } S_n = 1 + \frac{4}{5} + \frac{7}{5^2} + \frac{10}{5^3} + \dots + \frac{3n-2}{5^{n-1}}$$

$$\therefore \frac{1}{5} S_n = \frac{1}{5} + \frac{4}{5^2} + \frac{7}{5^3} + \dots + \frac{3n-5}{5^{n-1}} + \frac{3n-2}{5^n}$$

$$\therefore S_n - \frac{1}{5} S_n = 1 + \frac{3}{5} + \frac{3}{5^2} + \frac{3}{5^3} + \dots + \frac{3}{5^{n-1}} - \frac{3n-2}{5^n}$$

$$\Rightarrow \left(1 - \frac{1}{5}\right) S_n = 1 + \frac{3}{5} \left(1 + \frac{1}{5} + \frac{1}{5^2} + \dots + \frac{1}{5^{n-2}}\right) - \frac{3n-2}{5^n}$$

$$\Rightarrow \frac{4}{5} S_n = 1 + \frac{3}{5} \left\{ \frac{1 \left(1 - \frac{1}{5^{n-1}}\right)}{1 - \frac{1}{5}} \right\} - \frac{3n-2}{5^n}$$

$$\therefore S_n = \frac{a(1-r^n)}{1-r}, r < 1$$

$$= 1 + \frac{3}{5} \times \frac{5}{4} \left[1 - \frac{1}{5^{n-1}} \right] - \frac{3n-2}{5^n}$$

$$= 1 + \frac{3}{4} - \frac{3}{4 \cdot 5^{n-1}} - \frac{3n-2}{5^n} = \frac{7}{4} - \frac{3}{4 \cdot 5^{n-1}} - \frac{3n-2}{5^n}$$

$$\Rightarrow S_n = \frac{35}{16} - \left(\frac{15}{16 \times 5^{n-1}} + \frac{3n-2}{4 \cdot 5^{n-1}} \right)$$

$$= \frac{35}{16} - \left(\frac{15+12n-8}{16 \times 5^{n-1}} \right) = \frac{35}{16} - \frac{12n-7}{16 \times 5^{n-1}}$$

2. $1 + 2.2 + 3.2^2 + 4.2^3 + \dots + 100.2^{99}$ is

clearly an AGP with A.P. : $1 + 2 + 3 + \dots + 100$

and G.P. : $1 + 2 + 2^2 + \dots + 2^{99}$.

$$\text{Let } S = 1 + 2.2 + 3.2^2 + 4.2^3 + \dots + 100.2^{99}$$

$$\therefore 2S = 2 + 2.2^2 + 3.2^3 + \dots + 99.2^{99} + 100.2^{100}$$

$$\therefore S - 2S = (1 + 2 + 2^2 + 2^3 + \dots + 2^{99}) - 100.2^{100}$$

$$\Rightarrow -S = \frac{(2^{100} - 1)}{2 - 1} - 100.2^{100}$$

$$\left(\because S_n = \frac{a(r^n - 1)}{r - 1} \text{ and this is a G.P.} \right)$$

$$\text{with 100 terms, } a = 1, r = 2$$

$$= 2^{100} - 1 - 100.2^{100}$$

$$S = 2^{100} \cdot 100 - 2^{100} + 1 = 2^{100} \cdot (100 - 1) + 1$$

$$= 99.2^{100} + 1.$$

3. Let $S = 1 + \frac{2}{3} + \frac{6}{3^2} + \frac{10}{3^3} + \frac{14}{3^4} + \dots$ upto ∞

$$\frac{1}{3} S = \frac{1}{3} + \frac{2}{3^2} + \frac{6}{3^3} + \frac{10}{3^4} + \dots \text{ upto } \infty$$

$$\begin{aligned} \Rightarrow S - \frac{1}{3}S &= 1 + \frac{4}{3} + \frac{4}{3^2} + \frac{4}{3^3} + \frac{4}{3^4} + \dots \text{ upto } \infty \\ &= \frac{4}{3} + 4 \left[\frac{1}{3^2} + \frac{1}{3^3} + \frac{1}{3^4} + \dots \infty \right] \\ &= \frac{4}{3} + 4 \left[\frac{\frac{1}{3^2}}{1 - \frac{1}{3}} \right] \quad \left(\because S_{\infty} = \frac{a}{1-r} \right) \\ &= \frac{4}{3} + 4 \left[\frac{\frac{1}{9}}{\frac{2}{3}} \right] = \frac{4}{3} + \frac{4}{6} = \frac{12}{6} = 2. \end{aligned}$$

4. Let $S_{\infty} = 1 + 2(1 - \sin \theta) + 3(1 - \sin \theta)^2 + 4(1 - \sin \theta)^3 + \dots \infty$

$$\Rightarrow S_{\infty} = 1 + 2a + 3a^2 + 4a^3 + \dots \infty$$

(where $a = 1 - \sin \theta$)

$$\therefore a S_{\infty} = a + 2a^2 + 3a^3 + \dots \infty$$

$$\Rightarrow S_{\infty} - a S_{\infty} = 1 + a + a^2 + a^3 + \dots \infty$$

$$(1-a) S_{\infty} = \frac{1}{1-a}$$

$$\Rightarrow S_{\infty} = \frac{1}{(1-a)^2} = \frac{1}{(1-1+\sin \theta)^2}$$

$$= \frac{1}{\sin^2 \theta} = \operatorname{cosec}^2 \theta$$

$$\Rightarrow S_{\infty} = \operatorname{cosec}^2 \theta = 1 + \cot^2 \theta = 1 + (2\sqrt{2})^2 = 1 + 8 = 9.$$

5. Let $S = 4 + \frac{4+d}{5} + \frac{4+2d}{5^2} + \frac{4+3d}{5^3} + \dots \infty$

$$\therefore \frac{1}{5}S = \frac{4}{5} + \frac{4+d}{5^2} + \frac{4+2d}{5^3} + \dots \infty$$

$$\Rightarrow S - \frac{1}{5}S = 4 + \frac{d}{5} + \frac{d}{5^2} + \frac{d}{5^3} + \dots \infty$$

$$\Rightarrow \frac{4}{5}S = 4 + \frac{d}{5} \left[1 + \frac{1}{5} + \frac{1}{5^2} + \dots \infty \right]$$

$$\Rightarrow \frac{4}{5}S = 4 + \frac{d}{5} \times \frac{1}{1 - \frac{1}{5}} \Rightarrow \frac{4}{5}S = 4 + \frac{d}{5} \times \frac{5}{4}$$

$$\Rightarrow \frac{4}{5}S = 4 + \frac{d}{4} \Rightarrow S = 5 + \frac{5d}{16}$$

Given $S = 10$

$$\therefore 5 + \frac{5d}{16} = 10 \Rightarrow \frac{5d}{16} = 5 \Rightarrow d = 16.$$

SOME SPECIAL SERIES

KEY FACTS

1. Sum of first n natural numbers

Let $S = 1 + 2 + 3 + 4 + \dots + n$

Clearly, this is an A.P. with $a = 1, l = n, n = n$.

$$\therefore S = \frac{n}{2}(a+l) = \frac{n(n+1)}{2} \quad \text{or} \quad \Sigma n = \frac{n(n+1)}{2}, \text{ where } \Sigma \text{ stands for summation.}$$

2. Sum of the squares of first n natural numbers

Let $S = 1^2 + 2^2 + 3^2 + 4^2 + \dots + n^2$.

Then, $S = \Sigma n^2 = \frac{n(n+1)(2n+1)}{6}$

3. Sum of the cubes of first n natural numbers

Let $S = 1^3 + 2^3 + 3^3 + 4^3 + \dots + n^3$

Then, $S = \Sigma n^3 = \left\{ \frac{n(n+1)}{2} \right\}^2 = (\Sigma n)^2$

4. If $T_n = an^3 + bn^2 + cn + d$, then

$$S_n = a \Sigma n^3 + b \Sigma n^2 + c \Sigma n + dn$$

$$= a \left[\frac{n(n+1)}{2} \right]^2 + b \left[\frac{n(n+1)(2n+1)}{6} \right] + c \left[\frac{n(n+1)}{2} \right] + dn.$$

SOLVED EXAMPLES

Ex. 1. Find the n th term and then sum to n terms of the following series.

(i) $1^2 + (1^2 + 2^2) + (1^2 + 2^2 + 3^2) + \dots$ (ii) $3.5 + 4.7 + 5.9 + \dots$

Sol. (i) n th term of the given series

$$T_n = 1^2 + 2^2 + 3^2 + \dots + n^2 = \Sigma n^2 = \frac{n(n+1)(2n+1)}{6} = \frac{2n^3 + 3n^2 + n}{6} = \frac{n^3}{3} + \frac{n^2}{2} + \frac{n}{6}$$

$$\therefore S_n = \frac{1}{3} \cdot \Sigma n^3 + \frac{1}{2} \cdot \Sigma n^2 + \frac{1}{6} \cdot \Sigma n = \frac{1}{3} \cdot \frac{n^2(n+1)^2}{4} + \frac{1}{2} \cdot \frac{n(n+1)(2n+1)}{6} + \frac{1}{6} \cdot \frac{n(n+1)}{2}$$

$$= \frac{n(n+1)}{12} \{n(n+1) + (2n+1) + 1\} = \frac{n(n+1)(n^2 + 3n + 2)}{12} = \frac{n(n+1)(n+1)(n+2)}{12} = \frac{n(n+1)^2(n+2)}{12}$$

(ii) In the given series each term is the product of two factors. The factors 3, 4, 5, are in A.P. having 3 as the first term and 1 as the common difference, therefore the n th term of this A.P. = $3 + (n-1)1 = 2 + n$. Also, the factors 5, 7, 9, are in A.P. having 5 as the first term and 2 as the common difference, therefore the n th term of this A.P. = $5 + (n-1)2 = 2n + 3$

$$\therefore n\text{th term of the series} = (2+n)(2n+3)$$

$$\Rightarrow T_n = 2n^2 + 7n + 6$$

$$\therefore S_n = 2 \cdot \Sigma n^2 + 7 \Sigma n + 6n = \frac{2n(n+1)(2n+1)}{6} + \frac{7n(n+1)}{2} + 6n$$

$$= \frac{2n(n+1)(2n+1) + 21n(n+1) + 36n}{6} = \frac{4n^3 + 27n^2 + 59n}{6}$$

Ex. 2. What is the sum of the series $15^2 + 16^2 + 17^2 + \dots + 30^2$ equal to?

(Kerala PET 2004)

Sol. $15^2 + 16^2 + 17^2 + \dots + 30^2 = (1^2 + 2^2 + 3^2 + \dots + 30^2) - (1^2 + 2^2 + 3^2 + \dots + 14^2)$

$$= \sum_{k=1}^{30} k^2 - \sum_{k=1}^{14} k^2 = \frac{1}{6} \times 30 \times 31 \times (2 \times 30 + 1) - \frac{1}{6} \times 14 \times 15 \times (2 \times 14 + 1)$$

$$\left(\because \sum_{k=1}^n k^2 = \frac{1}{6} \times n(n+1)(2n+1) \right)$$

$$= \frac{30 \times 31 \times 61}{6} - \frac{14 \times 15 \times 29}{6} = 9455 - 1015 = \mathbf{8440}.$$

Ex. 3. What is the sum of the series $1^3 - 2^3 + 3^3 - 4^3 + \dots + 9^3$ equal to?

(AIEEE 2002)

Sol. Reqd. Sum = $(1^3 + 2^3 + 3^3 + 4^3 + \dots + 9^3) - 2(2^3 + 4^3 + 6^3 + 8^3)$

$$= (1^3 + 2^3 + 3^3 + \dots + 9^3) - 2^4(1^3 + 2^3 + 3^3 + 4^3)$$

$$= \sum_{k=1}^9 k^3 - 2^4 \sum_{k=1}^4 k^3 = \frac{9^2(9+1)^2}{4} - 2^4 \times \frac{4^2 \times (4+1)^2}{4}$$

$$\left(\because \sum_{k=1}^n k^3 = \frac{n^2(n+1)^2}{4} \right)$$

$$= \frac{81 \times 100}{4} - \frac{16 \times 16 \times 25}{4} = 2025 - 1600 = \mathbf{425}.$$

Ex. 4. Find the sum of n terms of the series whose n th term is $n^2 + 3^n$

Sol. $t_n = n^2 + 3^n$

$$\Rightarrow t_1 = 1^2 + 3^1$$

$$t_2 = 2^2 + 3^2$$

$$t_3 = 3^2 + 3^3$$

$$\begin{aligned} t_4 &= 4^2 + 3^4 \\ &\vdots \quad \vdots \\ t_n &= n^2 + 3^n \end{aligned}$$

∴ Adding columnwise, we get

$$\begin{aligned} t_1 + t_2 + t_3 + \dots + t_n &= (1^2 + 2^2 + 3^2 + \dots + n^2) + (3^1 + 3^2 + \dots + 3^n) \\ &= \sum_{k=1}^n k^2 + \frac{3 \cdot (3^n - 1)}{(3 - 1)} = \frac{n(n+1)(2n+1)}{6} + \frac{3}{2}(3^n - 1). \end{aligned}$$

Ex. 5. If S_1, S_2 and S_3 are the sums of the first n natural numbers, their squares, their cubes respectively, then show that $9S_2^2 = S_3(1 + 8S_1)$

Sol.

$$S_1 = \frac{n(n+1)}{2}, \quad S_2 = \frac{n(n+1)(2n+1)}{6}, \quad S_3 = \frac{n^2(n+1)^2}{4}$$

∴

$$\begin{aligned} 9S_2^2 &= 9 \left\{ \frac{1}{6} n(n+1)(2n+1) \right\}^2 = 9 \left\{ \frac{1}{36} n^2(n+1)^2(2n+1)^2 \right\} \\ &= \frac{1}{4} n^2(n+1)^2(4n^2 + 4n + 1) = \frac{1}{4} n^2(n+1)^2(4n(n+1) + 1) \\ &= \frac{1}{4} n^2(n+1)^2 \left(1 + 8 \left(\frac{1}{2} n(n+1) \right) \right) = S_3(1 + 8S_1). \end{aligned}$$

PRACTICE SHEET

1. What is the n th term of the series $1 + \frac{(1+2)}{2} + \frac{(1+2+3)}{3} + \dots$?

- (a) $\frac{n+1}{2}$ (b) $\frac{n(n+1)}{2}$
 (c) $n^2 - (n+1)$ (d) $\frac{(n+1)(2n+3)}{2}$

2. For $n \in N$, the sum of the series $2.3 + 3.5 + 4.7 + \dots + (n+1)(2n+1)$ is equal to

- (a) $\frac{n}{3}(2n^2 + 3n + 1)$ (b) $\frac{1}{6}n(n^2 + n - 1)$

- (c) $\frac{n}{3}(3n^2 + 5n + 11)$ (d) $\frac{n}{6}(4n^2 + 15n + 17)$

(AMU 2004)

3. The value of $1^2 - 2^2 + 3^2 - 4^2 + \dots + 11^2$ is equal to
 (a) 55 (b) 66 (c) 77 (d) 88

(Kerala PET 2011)

4. Sum of n terms of the series $1^3 + 3^3 + 5^3 + 7^3 + \dots$ is
 (a) $n^2(2n^2 - 1)$ (b) $2n^2 + 3n^2$
 (c) $n^3(n - 1)$ (d) $n^3 + 8n + 4$

(WBJEE 2010)

5. If $S_n = 1^3 + 2^3 + \dots + n^3$ and $T_n = 1 + 2 + 3 + \dots + n$, then
 (a) $S_n = T_n^2$ (b) $S_n = T_n^3$ (c) $S_n^2 = T_n$ (d) $S_3^2 = T_n$

(EAMCET 2007)

ANSWERS

1. (a) 2. (d) 3. (b) 4. (a) 5. (a)

HINTS AND SOLUTIONS

1. Reqd. n th term = $\frac{1+2+3+\dots+n}{n}$
 $= \frac{n(n+1)}{2} \times \frac{1}{n} = \frac{n+1}{2}$.

2. Let $S = 2 \cdot 3 + 3 \cdot 5 + 4 \cdot 7 + \dots + (n+1)(2n+1)$
 $= \sum_{k=1}^n (k+1)(2k+1) = \sum_{k=1}^n (2k^2 + 3k + 1)$
 $= 2 \sum_{k=1}^n k^2 + 3 \sum_{k=1}^n k + n$

$$\begin{aligned}
 &= 2 \times \frac{n(n+1)(2n+1)}{6} + 3 \times \frac{n(n+1)}{2} + n \\
 &= \frac{n}{6} [2(n+1)(2n+1) + 9(n+1) + 6] \\
 &= \frac{n}{6} [4n^2 + 6n + 2 + 9n + 9 + 6] \\
 &= \frac{n}{6} (4n^2 + 15n + 17).
 \end{aligned}$$

3. Let $S = 1^2 - 2^2 + 3^2 - 4^2 + \dots + 11^2$

$$\begin{aligned}
 &= (1^2 + 2^2 + 3^2 + 4^2 + \dots + 11^2) - 2(2^2 + 4^2 + 6^2 + 8^2 + 11^2) \\
 &= (1^2 + 2^2 + 3^2 + 4^2 + \dots + 11^2) - 2^3(1^2 + 2^2 + 3^2 + 4^2 + 5^2) \\
 &= \frac{11 \times (11+1) \times (2 \times 11 + 1)}{6} - \frac{8 \times 5 \times (5+1) \times (2 \times 5 + 1)}{6} \\
 &= \frac{11 \times 12 \times 23}{6} - \frac{8 \times 5 \times 6 \times 11}{6} = 22 \times 23 - 40 \times 11 \\
 &= 506 - 440 = \mathbf{66}.
 \end{aligned}$$

4. Let $S_n = 1^3 + 3^3 + 5^3 + 7^3 + \dots$ upto n terms
 n th term of the series $= (2n-1)^3$

$$\begin{aligned}
 \therefore S_n &= \sum_{k=1}^n (2k-1)^3 = \sum_{k=1}^n [8k^3 - 12k^2 + 6k] \\
 &= 8 \sum_{k=1}^n k^3 - 12 \sum_{k=1}^n k^2 + 6 \sum_{k=1}^n k - n \\
 &= 8 \frac{n^2(n+1)^2}{4} - 12 \times \frac{n(n+1)(2n+1)}{6} \\
 &\quad + 6 \times \frac{n(n+1)}{2} - n \\
 &= 2n^2(n^2 + 2n + 1) - 2n(2n^2 + 3n + 1) + 3n(n+1) - n \\
 &= 2n^4 + 4n^3 + 2n^2 - 4n^3 - 6n^2 - 2n + 3n^2 + 3n - n \\
 &= 2n^4 + n^2 = \mathbf{n^2(2n^2 - 1)}.
 \end{aligned}$$

5. $S_n = \sum n^3$

$$= \frac{n^2(n+1)^2}{4} = \left(\frac{n(n+1)}{2} \right)^2 \quad \dots(i)$$

$$T_n = \frac{n(n+1)}{2} \quad \dots(ii)$$

\therefore From (i) and (ii)

$$S_n = T_n^2$$

SELF ASSESSMENT SHEET

1. If the first term of an A.P. be 'a', second be 'b', and n th be '2a', then the sum of n terms is

(a) $\frac{3ab}{2(b-a)}$ (b) $\frac{2ab}{5(b-a)}$ (c) $\frac{ab}{2(b-a)}$ (d) $\frac{3ab}{(b-a)}$

(MPPET 2010)

2. If S_n denotes the sum of n terms of an A.P., then $S_{n+3} - 3S_{n+2} + 3S_{n+1} - S_n$ is equal to

(a) 0 (b) 1 (c) 2 (d) 3

(Kerala PET 2000)

3. The interior angles of a polygon are in A.P. The smallest angle is 120° and the common difference is 5° . The number of sides of the polygon is

(a) 8 (b) 9 (c) 12 (d) 19

(AMU 2010)

4. Divide 20 into four parts which are in A.P. and such that the product of the first and fourth is to the product of the second and third in the ratio 2 : 3. Find the product of the first and fourth term of the A.P.

(a) 12 (b) 16 (c) 20 (d) 25

5. The sum of the first p terms of an A.P. is q and the sum of the first q terms is p . Find the sum of the first $(p+q)$ terms.

(a) pq (b) $p-q$ (c) $-(p+q)$ (d) 0

6. In a G.P., the ratio of the sum of first 3 terms to that of the first 6 terms is 125 : 152. Find the common ratio of the G.P.

(a) $\frac{2}{3}$ (b) $\frac{3}{4}$ (c) $\frac{4}{3}$ (d) $\frac{3}{5}$

(J&K CET 2007)

7. The sum of n terms of the following series $1 + (1+x) + (1+x+x^2) + \dots$ will be

(a) $\frac{1-x^n}{1-x}$ (b) $\frac{x(1-x^n)}{1-x}$

(c) $\frac{n(1-x) - x(1-x^n)}{(1-x)^2}$ (d) None of the above

(AMU 2003)

8. If a_1, a_2, \dots, a_{50} are in G.P., then $\frac{a_1 - a_3 + a_5 - \dots + a_{49}}{a_2 - a_4 + a_6 - \dots + a_{50}}$ is equal to

(a) 0 (b) 1 (c) $\frac{a_1}{a_2}$ (d) $\frac{a_1}{a_{50}}$

(Kerala PET 2006)

9. If a, b, c are in G.P and x, y are the arithmetic means of a, b and b, c respectively, then $\frac{1}{x} + \frac{1}{y}$ is equal to

(a) $\frac{b}{2}$ (b) $\frac{b}{3}$ (c) $\frac{2}{b}$ (d) $\frac{3}{b}$

10. If S is the sum to infinity of a G.P. whose first term is 1, then the sum of its first n terms is

(a) $S \left(1 - \frac{1}{S}\right)^{n-1}$ (b) $S \left(1 - \frac{1}{S}\right)^n$

(c) $S \left\{1 - \left(1 - \frac{1}{S}\right)^{n-1}\right\}$ (d) $S \left\{1 - \left(1 - \frac{1}{S}\right)^n\right\}$

(AMU 2004)

11. If H is the harmonic mean between P and Q , then $\frac{H}{P} + \frac{H}{Q} =$

- (a) 2 (b) $\frac{1}{2}$ (c) $\frac{PQ}{P+Q}$ (d) $\frac{P+Q}{PQ}$

(VITEEE 2007)

12. If a, b, c are in G.P, then $\log_a 10, \log_b 10$ and $\log_c 10$ are in
(a) A.P (b) G.P (c) H.P. (d) None of these

13. If the sum of the roots of the quadratic equation $ax^2 + bx + c = 0$ be equal to the sum of the reciprocals of their squares, then bc^2, ca^2, ab^2 will be in
(a) A.P. (b) G.P. (c) H.P. (d) None of these

14. Let a_1, a_2, \dots, a_{10} be in A.P. and h_1, h_2, \dots, h_{10} be in H.P. If $a_1 = h_1 = 2$ and $a_{10} = h_{10} = 3$, then $a_4 h_7$ is
(a) 2 (b) 3 (c) 5 (d) 6 (IIT)

15. If a, b, c are in H.P, then $\frac{a}{b+c}, \frac{b}{c+a}, \frac{c}{a+b}$ will be in
(a) A.P. (b) G.P. (c) H.P. (d) None of these
(VITEEE 2011)

16. If $|x| < 1$, then the square root of the sum $1 + 2x + 3x^2 + 4x^3 + \dots \infty$
(a) $(1-x)$ (b) $(1+x)$ (c) $(1-x)^{-1}$ (d) $(1+x)^{-1}$
(Rajasthan PET 2007)

17. The sum to n terms of the series $1 + 2\left(1 + \frac{1}{n}\right) + 3\left(1 + \frac{1}{n}\right)^2 + \dots$ is
(a) n^2 (b) $n(n-1)$ (c) $(n+1)^2$ (d) $n(n+1)$
(EAMCET 2000)

18. Find the sum of n terms of the series whose n th term is $2n^2 + 3n$

- (a) $\frac{1}{2}n(n+1)(2n+5)$ (b) $\frac{1}{6}n(n+1)(4n+1)$
(c) $\frac{1}{3}(n^2 - 6n + 3)$ (d) $\frac{1}{6}n(n-1)(n^2 + 1)$

19. Find the sum of the series $1.2^2 + 3.3^2 + 5.4^2 + \dots$ to n terms

- (a) $\frac{n}{2}(n^3 + 4n^2 + 4n - 1)$ (b) $\frac{n}{2}(n+1)^2(2n+1)$
(c) $\frac{n^2}{3}(n+1)^2(2n+1)$ (d) $\frac{n}{3}(n^3 + 4n^2 - 2n + 1)$

20. If the sum of first n terms of an A.P. is cn^2 , then the sum of the squares of these n terms is

- (a) $\frac{n(4n^2 - 1)c^2}{6}$ (b) $\frac{n(4n^2 + 1)c^2}{6}$
(c) $\frac{n(4n^2 - 1)c^2}{3}$ (d) $\frac{n(4n^2 + 1)c^2}{3}$
(IIT 2009)

ANSWERS

1. (a) 2. (a) 3. (b) 4. (b) 5. (c) 6. (d) 7. (c) 8. (c) 9. (c) 10. (d)
11. (a) 12. (c) 13. (a) 14. (d) 15. (c) 16. (c) 17. (a) 18. (b) 19. (a) 20. (c)

HINTS AND SOLUTIONS

1. Let the common difference of the A.P. be ' d '. Then

$$d = b - a \quad \dots(i)$$

and $T_n = 2a = a + (n - 1)d$
 $\Rightarrow 2a = a + (n - 1)(b - a)$ (Using (i))

$$\Rightarrow 2a = a + bn - b - na + a$$

$$\Rightarrow b = n(b - a) \Rightarrow n = \frac{b}{b - a}$$

$$\therefore S_n = \frac{n}{2} \{a + T_n\} = \frac{b}{2(b - a)} (a + 2a) = \frac{3ab}{2(b - a)}$$

2. $S_{n+3} - 3S_{n+2} + 3S_{n+1} - S_n$
 $= (S_{n+3} - S_{n+2}) - 2(S_{n+2} - S_{n+1}) + (S_{n+1} - S_n)$
 $= t_{n+3} - 2t_{n+2} + t_{n+1} \quad \dots (i)$
 $(\because t_n = S_n - S_{n-1})$

$\therefore t_{n+1}, t_{n+2}, t_{n+3}$ are consecutive terms of an A.P,
 $2t_{n+2} = t_{n+1} + t_{n+3} \quad \dots(ii)$

\therefore From (i)

$$\text{Reqd. Sum} = (t_{n+3} + t_{n+1}) - 2t_{n+2} = 2t_{n+2} - 2t_{n+2} = 0 \quad \text{(Using (ii))}$$

3. Let the polygon have n sides.

Then, sum of its interior $\angle s$ (S_n) = $(2n - 4)$ rt. $\angle s$
 $= (n - 2) \times 180^\circ \quad \dots(i)$

Number of sides of the polygon = number of interior angles of the polygon = n .

The interior angles form an A.P. with first term = 120° and common difference 5° .

$$\therefore S_n = \frac{n}{2} [2 \times 120^\circ + (n - 1) \times 5^\circ]$$

$$= \frac{n}{2} [240^\circ + 5n - 5^\circ] \quad \dots(ii)$$

From (i) and (ii),

$$(n - 2) \times 180^\circ = \frac{n}{2} [240^\circ + 5n - 5^\circ]$$

$$\begin{aligned} \Rightarrow (n-2) \times 360 &= 5n^2 + 235n \\ \Rightarrow 5n^2 + 235n - 360n + 720 &= 0 \\ \Rightarrow 5n^2 - 125n + 720 &= 0 \Rightarrow n^2 - 25n + 144 = 0 \\ \Rightarrow (n-16)(n-9) &= 0 \Rightarrow n = 16 \text{ or } 9. \\ \text{when } n = 16, \text{ the last angle } a_n &= a + (n-1)d \\ &= 120^\circ + (16-1) \times 5^\circ = 195^\circ \end{aligned}$$

which is not possible.

Hence, $n = 9$.

4. Let the required four parts be $(a - 3d)$, $(a - d)$, $(a + d)$, $(a + 3d)$

Given, $a - 3d + a - d + a + d + a + 3d = 20$

$$\Rightarrow 4a = 20 \Rightarrow a = 5$$

Also, $\frac{(a - 3d)(a + 3d)}{(a - d)(a + d)} = \frac{2}{3}$

$$\Rightarrow 3(a^2 - 9d^2) = 2(a^2 - d^2)$$

$$\Rightarrow 3(25 - 9d^2) = 2(25 - d^2)$$

$$\Rightarrow 75 - 27d^2 = 50 - 2d^2 \Rightarrow 25 = 25d^2$$

$$\Rightarrow d = \pm 1$$

Taking $d = 1$, the numbers are $5 - 3, 5 - 1, 5 + 1, 5 + 3$, i.e., $2, 4, 6, 8$

Taking $d = -1$, the numbers are $5 + 3, 5 + 1, 5 - 1, 5 - 3$, i.e., $8, 6, 4, 2$

\therefore Required product in either case $= 2 \times 8 = 16$.

5. Since $S_n = \frac{n}{2}[2a + (n-1)d]$ for an A.P. whose first term

$= a$, common difference $= d$, number of terms $= n$.

$$\therefore S_p = q = \frac{p}{2}(2a + (p-1)d)$$

$$\Rightarrow 2q = 2ap + p(p-1)d \quad \dots(i)$$

$$S_q = p = \frac{q}{2}(2a + (q-1)d)$$

$$\Rightarrow 2p = 2aq + q(q-1)d \quad \dots(ii)$$

Subtracting eqn (ii) from eqn (i), we get

$$2(q-p) = 2a(p-q) + (p^2 - q^2)d - (p-q)d$$

$$\Rightarrow -2(p-q) = 2a(p-q) + (p-q)(p+q)d - (p-q)d$$

$$-2 = 2a + [(p+q)-1]d \quad \dots(iii)$$

Now $S_{p+q} = \frac{p+q}{2}[2a + (p+q-1)d]$

$$= \frac{p+q}{2} \times -2 \quad \text{(From (iii))}$$

$$= -(p+q).$$

6. Let a be the first term and r the common ratio of the G.P.

Then, $\frac{S_3}{S_6} = \frac{125}{152} \Rightarrow \frac{a(r^3 - 1)}{a(r^6 - 1)} = \frac{125}{152}$

$$\frac{(r^3 - 1)}{(r^6 - 1)} = \frac{125}{152} \Rightarrow \frac{(r^3 - 1)}{(r^3 + 1)(r^3 - 1)} = \frac{125}{152}$$

$$\Rightarrow r^3 + 1 = \frac{152}{125} \Rightarrow r^3 = \frac{152}{125} - 1 = \frac{27}{125}$$

$$\Rightarrow r^3 = \left(\frac{3}{5}\right)^3 \Rightarrow r = \frac{3}{5}.$$

7. $S_n = 1 + (1+x) + (1+x+x^2) + \dots$ n terms

$$\Rightarrow S_n = \frac{1}{1-x} [(1-x) + (1-x)(1+x) + (1-x)(1+x+x^2) + \dots \text{ to } n \text{ terms}]$$

$$= \frac{1}{1-x} [(1-x) + (1-x^2) + (1-x^3) + \dots \text{ to } n \text{ terms}]$$

$$= \frac{1}{1-x} [n - (x + x^2 + x^4 + \dots \text{ to } n \text{ terms})]$$

$$= \frac{1}{1-x} \left[n - \frac{x(1-x^n)}{1-x} \right] \quad \left(\because S_n = \frac{a(1-r^n)}{1-r} \right)$$

$$= \frac{n(1-x) - x(1-x^n)}{(1-x)^2}.$$

8. Let the first term of the G.P. be a and common ratio r . Then,

$$\frac{a_1 - a_3 + a_5 - \dots + a_{49}}{a_2 - a_4 + a_6 - \dots + a_{50}} = \frac{a - ar^2 + ar^4 - \dots + ar^{48}}{ar - ar^3 + ar^5 - \dots + ar^{49}}$$

$$= \frac{a(1 - r^2 + r^4 - \dots + r^{48})}{ar(1 - r^2 + r^4 - \dots + r^{48})}$$

$$= \frac{a}{ar} = \frac{a_1}{a_2}.$$

9. a, b, c are in G.P $\Rightarrow b^2 = ac \quad \dots(i)$

x is the A.M. of $a, b \Rightarrow x = \frac{a+b}{2} \quad \dots(ii)$

y is the A.M. of $b, c \Rightarrow y = \frac{b+c}{2} \quad \dots(iii)$

Now $\frac{1}{x} + \frac{1}{y} = \frac{2}{a+b} + \frac{2}{b+c} = \frac{2(b+c) + 2(a+b)}{(a+b)(b+c)}$

$$= \frac{2(a+2b+c)}{ab+b^2+ac+bc} = \frac{2(a+2b+c)}{ab+b^2+b^2+bc}$$

$$= \frac{2(a+2b+c)}{b(a+2b+c)} = \frac{2}{b}.$$

10. Let the first term of the G.P. be a and common ratio r .

Given $S = \frac{a}{1-r} \Rightarrow S = \frac{1}{1-r}$

$$\Rightarrow 1-r = \frac{1}{S} \Rightarrow r = 1 - \frac{1}{S}$$

Let S_n be the sum of first n terms of the series. Then,

$$S_n = \frac{1(1-r^n)}{1-r} \quad (\because |r| < 1)$$

$$= \frac{1 \left(1 - \left(1 - \frac{1}{S} \right)^n \right)}{\left(1 - \left(1 - \frac{1}{S} \right) \right)} = S \left\{ 1 - \left(1 - \frac{1}{S} \right)^n \right\}.$$

11. H being the harmonic mean between P and Q .

$$H = \frac{2PQ}{P+Q}$$

$$\therefore \frac{H}{P} + \frac{H}{Q} = \frac{2Q}{P+Q} + \frac{2P}{P+Q} = \frac{2(P+Q)}{P+Q} = 2.$$

12. a, b, c are in G.P. $\Rightarrow b^2 = ac$

$$\begin{aligned} \Rightarrow \frac{1}{\log_a 10} + \frac{1}{\log_c 10} &= \log_{10} a + \log_{10} c = \log_{10} (ac) \\ &= \log_{10} b^2 = 2 \log_{10} b = \frac{2}{\log_b 10} \end{aligned}$$

$$\Rightarrow \frac{1}{\log_a 10}, \frac{1}{\log_b 10}, \frac{1}{\log_c 10} \text{ are in A.P.}$$

$$\Rightarrow \log_a 10, \log_b 10, \log_c 10 \text{ are in H.P.}$$

13. Let α, β , be the roots of the equation $ax^2 + bx + c = 0$. Then

$$\text{Sum of roots} = \alpha + \beta = -\frac{b}{a} \quad \dots(i)$$

$$\text{Product of roots} = \alpha\beta = \frac{c}{a} \quad \dots(ii)$$

$$\text{Given, } \alpha + \beta = \frac{1}{\alpha^2} + \frac{1}{\beta^2}$$

$$\Rightarrow \alpha + \beta = \frac{\alpha^2 + \beta^2}{\alpha^2\beta^2} = \frac{(\alpha + \beta)^2 - 2\alpha\beta}{\alpha^2\beta^2}$$

$$\left(-\frac{b}{a}\right) = \frac{\left(-\frac{b}{a}\right)^2 - \frac{2c}{a}}{\left(\frac{c}{a}\right)^2} \quad (\text{From (i) and (ii)})$$

$$-\frac{b}{a} = \frac{\frac{b^2}{a^2} - \frac{2c}{a}}{\frac{c^2}{a^2}} \Rightarrow \frac{-bc^2}{a^3} = \frac{b^2 - 2ac}{a^2}$$

$$\Rightarrow -\frac{bc^2}{a} = b^2 - 2ac \Rightarrow -bc^2 = ab^2 - 2ac$$

$$\Rightarrow 2a^2c = ab^2 + bc^2$$

$$\Rightarrow ab^2, a^2c, bc^2 \text{ are in A.P.}$$

14. Let d be the common difference of the given A.P. a_1, a_2, \dots, a_{10} .

$$\text{Then, given, } a_1 = 2 \text{ and } a_{10} = a_1 + 9d = 3$$

$$\Rightarrow 2 + 9d = 3 \Rightarrow d = \frac{1}{9}$$

$$\Rightarrow a_4 = a_1 + 3d = 2 + 3 \times \frac{1}{9} = 2\frac{1}{3} = \frac{7}{3} \quad \dots (i)$$

Now, h_1, h_2, \dots, h_{10} are in H.P.

$$\Rightarrow \frac{1}{h_1}, \frac{1}{h_2}, \dots, \frac{1}{h_{10}} \text{ are in A.P.}$$

Let d_1 be the common difference of this A.P.

$$\text{Given, } h_1 = 2 \Rightarrow \text{First term of the A.P.} = \frac{1}{h_1} = \frac{1}{2}$$

$$\text{Also, } h_{10} = 3 \Rightarrow \frac{1}{h_{10}} = \frac{1}{h_1} + 9d_1$$

$$\Rightarrow \frac{1}{3} = \frac{1}{2} + 9d_1 \Rightarrow 9d_1 = \frac{1}{3} - \frac{1}{2} = -\frac{1}{6}$$

$$\Rightarrow d_1 = -\frac{1}{54}$$

$$\therefore \frac{1}{h_7} = \frac{1}{h_1} + 6d_1 = \frac{1}{2} + 6 \times -\frac{1}{54} = \frac{1}{2} - \frac{1}{9} = \frac{7}{18}$$

$$\Rightarrow h_7 = \frac{18}{7} \quad \dots(ii)$$

$$\therefore \text{From (i) and (ii) } a_4 h_7 = \frac{7}{3} \times \frac{18}{7} = 6.$$

15. a, b, c are in H.P.

$$\Rightarrow \frac{1}{a}, \frac{1}{b}, \frac{1}{c} \text{ are in A.P.}$$

$$\Rightarrow \frac{a+b+c}{a}, \frac{a+b+c}{b}, \frac{a+b+c}{c} \text{ are in A.P.}$$

(Multiplying each term by $(a+b+c)$)

$$\Rightarrow 1 + \frac{b+c}{a}, 1 + \frac{a+c}{b}, 1 + \frac{a+b}{c} \text{ are in A.P.}$$

$$\Rightarrow \frac{b+c}{a}, \frac{a+c}{b}, \frac{a+b}{c} \text{ are in A.P.}$$

(Subtracting 1 from each term)

$$\Rightarrow \frac{a}{b+c}, \frac{b}{a+c}, \frac{c}{a+b} \text{ are in H.P.}$$

16. Let $S_\infty = 1 + 2x + 3x^2 + 4x^3 + \dots \infty \quad \dots(i)$

S_∞ is an infinite A.G.P. with A.P. : $1 + 2 + 3 + \dots \infty$

and G.P. : $1 + x + x^2 + x^3 + \dots \infty$

The common ratio of the A.G.P. is x

$$\therefore x S_\infty = x + 2x^2 + 3x^3 + \dots \infty \quad \dots(iii)$$

Eq (i) - Eq (ii)

$$\Rightarrow (1-x) S_\infty = 1 + x + x^2 + x^3 + \dots \infty$$

$$\Rightarrow (1-x) S_\infty = \frac{1}{1-x} \quad \left(\because S_\infty = \frac{a}{1-r} \right)$$

$$\Rightarrow S_\infty = \frac{1}{(1-x)^2} = (1-x)^{-2}$$

$$\therefore \text{Square root of } S_\infty = (1-x)^{-1}.$$

$$17. S_n = 1 + 2\left(1 + \frac{1}{n}\right) + 3\left(1 + \frac{1}{n}\right)^2 + \dots + n\left(1 + \frac{1}{n}\right)^{n-1}$$

$$\therefore \left(1 + \frac{1}{n}\right) S_n = \left(1 + \frac{1}{n}\right) + 2\left(1 + \frac{1}{n}\right)^2 + \dots$$

$$+ (n-1)\left(1 + \frac{1}{n}\right)^{n-1} + n\left(1 + \frac{1}{n}\right)^n$$

$$\left(\because S_n \text{ is an A.G.P with common ratio } \left(1 + \frac{1}{n}\right) \right)$$

$$\begin{aligned} \Rightarrow S_n \left[1 - \left(1 + \frac{1}{n} \right) \right] &= 1 + \left(1 + \frac{1}{n} \right) + \left(1 + \frac{1}{n} \right)^2 + \dots \\ &\quad \left(1 + \frac{1}{n} \right)^{n-1} - n \left(1 + \frac{1}{n} \right)^n \\ \Rightarrow -\frac{1}{n} S_n &= \frac{1 \left(\left(1 + \frac{1}{n} \right)^n - 1 \right)}{\left(1 + \frac{1}{n} \right) - 1} - n \left(1 + \frac{1}{n} \right)^n \\ \Rightarrow -\frac{1}{n} S_n &= n \left[\left(1 + \frac{1}{n} \right)^n - 1 \right] - n \left(1 + \frac{1}{n} \right)^n \\ \Rightarrow -\frac{1}{n} S_n &= -n \Rightarrow S_n = n^2. \end{aligned}$$

$$\begin{aligned} \mathbf{18.} \quad T_n = 2n^2 + 3n \quad \therefore S_n &= 2 \sum_{k=1}^n k^2 + 3 \sum_{k=1}^n k \\ &= 2 \times \frac{1}{6} \times n(n+1)(2n+1) + 3 \times \frac{n(n+1)}{2} \\ &= \frac{1}{3} \times n(n+1)(2n+1) + \frac{3}{2} n(n+1) \\ &= n(n+1) \left[\frac{2n+1}{3} + \frac{3}{2} \right] = n(n+1) \left[\frac{4n+2+9}{6} \right] \\ &= \frac{n(n+1)(4n+1)}{6}. \end{aligned}$$

19. n th term (T_n) of the given series

$$\begin{aligned} &= (2n-1)(n+1)^2 = (2n-1)(n^2+2n+1) \\ &= 2n^3 - n^2 + 4n^2 - 2n + 2n - 1 = 2n^3 + 3n^2 - 1 \\ \therefore S_n &= 2 \sum_{n=1}^n k^3 + 3 \sum_{k=1}^n k^2 - n \end{aligned}$$

$$\begin{aligned} &= 2 \times \frac{n^2(n+1)^2}{4} + 3 \times \frac{1}{6} \times n(n+1)(2n+1) - n \\ &= \frac{1}{2} [n^2(n^2+2n+1) + (n^2+n)(2n+1) - 2n] \\ &= \frac{1}{2} [n^4 + 2n^3 + n^2 + 2n^3 + 2n^2 + n^2 + n - 2n] \\ &= \frac{1}{2} [n^4 + 4n^3 + 4n^2 - n] = \frac{n}{2} [n^3 + 4n^2 + 4n - 1]. \end{aligned}$$

20. Let the sum of first n terms of the A.P, $S_n = cn^2$

$$\therefore S_{n-1} = c(n-1)^2 = c(n^2 - 2n + 1)$$

$$\begin{aligned} \therefore n\text{th term of the A.P.} &= S_n - S_{n-1} \\ &= cn^2 - c(n^2 - 2n + 1) = c(2n - 1) \end{aligned}$$

Let S_n^2 be the sum of the squares of these n terms. Then,

$$\begin{aligned} S_n^2 &= \sum_{k=1}^n t_k^2 \quad \text{where } t_k = c(2k-1) \\ &= \sum_{k=1}^n [c^2(2k-1)^2] = \sum_{k=1}^n [c^2 \cdot 4k^2 - c^2 \cdot 4k + c^2] \\ &= 4c^2 \sum_{k=1}^n k^2 - 4c^2 \sum_{k=1}^n k + c^2 n \\ &= 4c^2 \times \frac{n(n+1)(2n+1)}{6} - 4c^2 \times \frac{n(n+1)}{2} + c^2 n \\ &= \frac{c^2 n}{6} [4(n+1)(2n+1) - 12(n+1) + 6] \\ &= \frac{c^2 n}{6} [8n^2 + 12n + 4 - 12n - 12 + 6] \\ &= \frac{c^2 n}{6} [8n^2 - 2] = \frac{n(4n^2 - 1)c^2}{3}. \end{aligned}$$

SELF ASSESSMENT SHEET

1. If $A = \begin{bmatrix} 4 & 3 \\ 2 & 5 \end{bmatrix}$, find x and y such that $A^2 - xA + yI = 0$

- (a) $x = 9, y = -14$ (b) $x = 14, y = 9$
 (c) $x = 9, y = 14$ (d) $x = -9, y = 14$

2. If A and B are two square matrices of the same order such that $AB = B$ and $BA = A$, then $A^2 + B^2$ is always equal to
 (a) I (b) $A + B$ (c) $2AB$ (d) $2BA$

3. If $\begin{bmatrix} a & -1 & 5 \\ b & 3 & -4 \end{bmatrix} \begin{bmatrix} 4 & 3 \\ 1 & 0 \\ 7 & -6 \end{bmatrix} = \begin{bmatrix} 34 & -30 \\ -1 & 42 \end{bmatrix}$, then (a, b) is equal to

- (a) $(1, 3)$ (b) $(-2, 4)$ (c) $(0, 6)$ (d) $(2, -3)$

4. If $A = \begin{bmatrix} 0 & -3 & -4/3 \\ 3 & 0 & -1/4 \\ 4/3 & 1/4 & 0 \end{bmatrix}$, then $\det. (A + A^T)$ is equal to

- (a) 0 (b) 1 (c) 2 (d) 3

5. If every element of a determinant of order 3 of value Δ is multiplied by 5, then the value of the new determinant is
 (a) Δ (b) 5Δ (c) 25Δ (d) 125Δ

6. If $M = \begin{vmatrix} a & l & p \\ b & m & q \\ c & n & r \end{vmatrix}$ and $N = \begin{vmatrix} p & q & r \\ a & b & c \\ l & m & n \end{vmatrix}$, then

- (a) $M' = N$ (b) $M = -N$ (c) $M = N^2$ (d) $M = N^3$

7. If $\begin{vmatrix} -a^2 & ab & ac \\ ab & -b^2 & bc \\ ac & bc & -c^2 \end{vmatrix} = k a^2 b^2 c^2$, then k is equal to

- (a) -4 (b) 2 (c) 4 (d) 8

8. The matrix $\begin{bmatrix} \lambda & -1 & 4 \\ -3 & 0 & 1 \\ -1 & 1 & 2 \end{bmatrix}$ is invertible if

- (a) $\lambda \neq -20$ (b) $\lambda \neq -19$ (c) $\lambda \neq -18$ (d) $\lambda \neq -17$

9. The element in the first row and third column of the inverse of the matrix $\begin{bmatrix} 1 & 2 & -3 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{bmatrix}$ is

- (a) -2 (b) 0 (c) 1 (d) 7

10. The system of linear equations $2x + 2y = 5$, $5x + ky = 9$ has a unique solution if

- (a) $k \neq 5$ (b) $k = 0$ (c) $k \neq -1$ (d) $k \neq 2$

11. The system of equations $3x + 4y = 2$ and $6x + 8y = 4$ has
 (a) No solution (b) a unique solution
 (c) two distinct solutions (d) infinitely many solutions.

12. The system of equations $x + 3y = 5$, $2x + 6y = 8$ has
 (a) No solution (b) a unique solution
 (c) infinitely many solutions (d) None of these

ANSWERS

1. (c) 2. (b) 3. (c) 4. (a) 5. (d) 6. (a) 7. (c) 8. (d) 9. (d) 10. (a)
 11. (d) 12. (a)

HINTS AND SOLUTIONS

1. $A = \begin{bmatrix} 4 & 3 \\ 2 & 5 \end{bmatrix} \Rightarrow A^2 = A \cdot A = \begin{bmatrix} 4 & 3 \\ 2 & 5 \end{bmatrix} \begin{bmatrix} 4 & 3 \\ 2 & 5 \end{bmatrix}$
 $= \begin{bmatrix} 16+6 & 12+15 \\ 8+10 & 6+25 \end{bmatrix} = \begin{bmatrix} 22 & 27 \\ 18 & 31 \end{bmatrix}$

Given $A^2 - xA + yI = 0 \Rightarrow A^2 - xA = -yI$

$\Rightarrow \begin{bmatrix} 22 & 27 \\ 18 & 31 \end{bmatrix} - x \begin{bmatrix} 4 & 3 \\ 2 & 5 \end{bmatrix} = -y \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

$\Rightarrow \begin{bmatrix} 22-4x & 27-3x \\ 18-2x & 31-5x \end{bmatrix} = \begin{bmatrix} -y & 0 \\ 0 & -y \end{bmatrix}$

$\Rightarrow 22 - 4x = -y, 27 - 3x = 0$

Now $27 - 3x = 0 \Rightarrow 3x = 27 \Rightarrow x = 9$

$\therefore 22 - 4x = -y \Rightarrow 22 - 36 = -y$

$\Rightarrow -y = -14 \Rightarrow y = 14.$

2. We know that if $AB = B$ and $BA = A$, then A and B are idempotent matrices, i.e., $A^2 = A$ and $B^2 = B$.

$A^2 + B^2 = A + B$

Alternatively,

$A^2 + B^2 = A \cdot A + B \cdot B = A \cdot (BA) + B \cdot (AB)$

$= (AB)A + (BA)B$

(Matrix multiplication is associative)

$= BA + AB$ ($\because AB = B, BA = A$)

$= A + B$ ($\because BA = A, AB = B$)

3. $\begin{bmatrix} a & -1 & 5 \\ b & 3 & -4 \end{bmatrix}_{2 \times 3} \begin{bmatrix} 4 & 3 \\ 1 & 0 \\ 7 & -6 \end{bmatrix}_{3 \times 2} = \begin{bmatrix} 34 & -30 \\ -1 & 42 \end{bmatrix}_{2 \times 2}$

LHS = $\begin{bmatrix} 4a - 1 + 35 & 3a + 0 - 30 \\ 4b + 3 - 28 & 3b + 0 + 24 \end{bmatrix} = \begin{bmatrix} 4a + 34 & 3a - 30 \\ 4b - 25 & 3b + 24 \end{bmatrix}$

RHS = $\begin{bmatrix} 34 & -30 \\ -1 & 42 \end{bmatrix}$

Equating corresponding elements we have

$4a + 34 = 34 \Rightarrow a = 0$ and $4b - 25 = -1 \Rightarrow 4b = 24 \Rightarrow b = 6$

$3a - 30 = -30 \Rightarrow a = 0$ and $3b + 24 = 42 \Rightarrow 3b = 18 \Rightarrow b = 6$

$\therefore a = 0, b = 6.$

WHY MODULUS JUNIOR COLLEGE ?

Best IITJEE Faculty : Team led by an Alumni of IIT Delhi, **MODULUS** has expert faculty to provide IITJEE coaching, mentoring and counselling to the students to reach their goal.

Exceptional Teaching Methodology : Concept Oriented Teaching rather than Rote Learning. Sufficient self-study time is given to grasp the concepts and to apply them to solve the difficult problems. Classes will be interactive to keep students curious and interested in subject.

“I listen - I forget,
I read -I remember,
I do - I understand”

following this saying, measures like asking students to teach some subtopic will be taken.

Scientifically Researched Program : IITJEE experts /toppers recommend for every one hour of teaching, one hour of self study should be given. Triumph program is designed keeping this in mind.

Optimum Batch Size : Batch size of 30-35 students to provide individual attention to each and every student.

Excellent Study material : Complete, well structured and comprehensive study material prepared by our Subject Matter Experts will be provided.

Regular Exams : IITJEE(JEE MAINS/ADVANCE), BITSAT & EAMCET pattern exams will be conducted weekly to test students followed by test analysis to diagnose the students weak areas.

Students Mentorship Program : Students will be assigned Mentors to resolve their day to day problems. MENTORS will follow up with students on regular basis to monitor their progress and will help them to stay focused on their goal.

Overall Personality Development : Regular sessions of YOGA, MEDITATION, SPORTS, Moral and Ethical Behavior will be conducted for students holistic development.

Regular PTM & Motivational seminars: Regular Parent - Teacher meetings will be conducted to discuss the performance of their ward. Regular motivational seminars by IIT Alumni , IIT JEE Experts to instigate the deep desire in students to work hard to get admissions into world class institutes.

Career Guidance to students after completion of their Intermediate course.

Transport Facility: Transport Facility will be provided to needy students.

Hostel Facility: Separate hostel facility for BOYS and GIRLS will be provided.

MODULUS Junior College

WEEK DAYS / WEEKENDS

Class Room Programs

7th

For 7th Class Students
ENRICH TWO YEAR

8th

For 8th Class Students
ENRICH ONE YEAR

The main aim of the programme is to enrich logical thinking, IQ besides laying strong foundation in **MATHS, PHYSICS, CHEMISTRY**.
This program also helps students to excel in school exams.

9th

For 9th Class Students
REFINE TWO YEAR

10th

For 10th Class Students
REFINE ONE YEAR

The goal of this program is to lay strong foundation for the most prestigious competitive exams like **NTSE, IIT-JEE, KVPY, NSEJS** and other Olympiads.
This also helps students to perform exceptionally in board exams.

11th

For 11th Class Students
TRIUMPH TWO YEAR

12th

For 12th Class Students
TRIUMPH ONE YEAR

The above programs will help the student to perform well in IIT-JEE Mains and Advanced exams. This also equip the student excel in other engineering and board exams.

ADDRESS

OPP. ZENAS INTERNATIONAL SCHOOL , Near Miyapur Metro First Pillar ,
Mathrusri Nagar , Miyapur ,Hyderabad.

